

The Use of Jejunal Tube Feeding in Children: A Position Paper by the Gastroenterology and Nutrition Committees of the European Society for Paediatric Gastroenterology, Hepatology, and Nutrition 2019

Ilse J. Broekaert, †Jackie Falconer, ‡Jiri Bronsky, §Frederic Gottrand, ||Luigi Dall'Oglio, ¶Eunice Goto, *Iva Hojsak, ††Jessie Hulst, ††Brigitte Kochavi, §§Alexandra Papadopoulou, ||||Carmen Ribes-Koninckx, ¶¶Michela Schaeppi, ##Steven Werlin, ***Michael Wilschanski, and †††Nikhil Thapar*

ABSTRACT

Objectives: Jejunal tube feeding (JTF) is increasingly becoming the standard of care for children in whom gastric tube feeding is insufficient to achieve caloric needs. Given a lack of a systematic approach to the care of JTF in paediatric patients, the aim of this position paper is to provide expert guidance regarding the indications for its use and practical considerations to optimize its utility and safety.

Methods: A group of members of the Gastroenterology and Nutrition Committees of the European Society of Paediatric Gastroenterology Hepatology and Nutrition and of invited experts in the field was formed in September 2016 to produce this clinical guide. Seventeen clinical questions treating indications and contraindications, investigations before placement, techniques of placement, suitable feeds and feeding regimen, weaning from JTF, complications, long-term care, and ethical considerations were addressed. A systematic literature search was performed from 1982 to November 2018 using PubMed, the MEDLINE, and Cochrane Database of Systematic Reviews. Grading of Recommendations, Assessment, Development, and Evaluation was applied to evaluate the outcomes.

During a consensus meeting, all recommendations were discussed and finalized. In the absence of evidence from randomized controlled trials, recommendations reflect the expert opinion of the authors.

Results: A total of 33 recommendations were voted on using the nominal voting technique.

Conclusions: JTF is a safe and effective means of enteral feeding when gastric feeding is insufficient to meet caloric needs or is not possible. The decision to place a jejunal tube has to be made by close cooperation of a multidisciplinary team providing active follow-up and care.

Key Words: children, clinical guide, feeding, jejunal tube, multidisciplinary team, recommendations

(JPGN 2019;69: 239–258)

What Is Known

- Jejunal tube feeding is increasingly becoming the standard of care for children in whom gastric tube feeding is insufficient to achieve caloric needs.
- There is a lack of expert guidance regarding the indications and practical considerations to optimize its utility and safety in clinical practice.

What Is New

- Jejunal tube feeding is a safe and effective means of enteral feeding when gastric feeding is insufficient to meet caloric needs.
- The decision to place a jejunal tube has to be made by a multidisciplinary team, working in close cooperation and providing active follow-up and care.

Jejunal tube feeding (JTF) is defined as postpyloric feeding through a feeding tube with its tip placed at least 40 cm distally to ligament of Treitz. JTF bypasses the stomach when gastric feeding is not tolerated or associated with unacceptable complications including significant gastroesophageal reflux disease (GERD).

There is growing evidence suggesting the increased use of JTF in children (1–3) with a number of recent recommendations suggesting that feeding by jejunal tube (JT) is a valid option in infants or children who fail intragastric feeding (4–9). In parallel, a

Received January 7, 2019; accepted April 25, 2019.

From the *Faculty of Medicine, University Children's Hospital, University of Cologne, Cologne, Germany, the †Paediatric Gastroenterology and Nutrition unit, Chelsea and Westminster Hospital, Fulham, London, UK, the ‡Department of Paediatrics, University Hospital Motol, Prague, Czech Republic, the §Department of Paediatric Gastroenterology, Hepatology and Nutrition, CHU Lille, University Lille, Lille, France, the ||Digestive Endoscopy and Surgical Unit, Bambino Gesù Children's Hospital, Rome, Italy, the ¶Paediatric Gastroenterology and Nutrition Unit, Chelsea and Westminster Hospital, Fulham, London, UK, the

#Children's Hospital Zagreb, University of Zagreb School of Medicine, Zagreb, the **School of Medicine, University J.J. Strossmayer, Osijek, Croatia, the ††Department of Paediatric Gastroenterology, Erasmus Medical Centre – Sophia Children's Hospital, Rotterdam, The Netherlands, the †††Paediatric Gastroenterology Unit, The Edmond and Lily Safra Children's Hospital, The Haim Sheba Medical Center, Ramat Gan, Israel, and the §§Division of Gastroenterology, Hepatology and Nutrition, First Department of Pediatrics, University of Athens, Children's Hospital "Agia Sofia", Athens, Greece

number of these and other articles also report on the safety, efficacy, and limitations associated with it (10–12). Retrospective studies show that gastrojejunal tube (GJT) feeding is a safe method to improve nutritional status; however, because of the frequent need for tube maintenance and replacement leading to increased morbidity, GJT feeding is rather a transitory alternative to, for example, surgical Roux-en-Y jejunostomy or antireflux surgery (10–12).

To our knowledge, there is little clear guidance as to the indications for the use of JTF or practical aspects related to its utility in clinical management. This article seeks to address some of these issues.

A number of factors should be considered, however, before placement of a JT, or indeed a GJT. The symptoms of feeding failure such as nausea, vomiting, gagging, retching, and volume intolerance may be caused by anatomical or, indeed, nongastrointestinal problems, which will need to be dealt with before considering placement of a JT.

The management of a child awaiting a jejunal feeding tube should begin well before its insertion and involve a multidisciplinary team (MDT) of health care providers who are familiar with, and have access to, a range of alternative strategies to the insertion of such a feeding tube. These may include feed or regimen changes, specific feeding therapy, speech and swallow assessments, and access to psychological support. The MDT should, arguably, include a paediatric gastroenterologist, nurse, psychologist, dietician, and a speech and language therapist.

Adequate planning, including discussion of ethical issues, warrants that all parties have a clear understanding of the indication and rationale for placement of a JT. In addition, ongoing and future strategies to increase possible oral feeding and enable weaning off the JT should be discussed.

The aim of this European Society for Paediatric Gastroenterology, Hepatology, and Nutrition (ESPGHAN) position statement is to provide a comprehensive guide for health care professionals on the safe, effective, and appropriate use of jejunal feeding tubes in children and young adults.

METHODOLOGY

Under the auspices of ESPGHAN, a working group (WG) consisting of members from the gastrointestinal (GI) and Nutrition Committees and experts in the field, including paediatric gastroenterologists, dieticians, a nurse, and a paediatric surgeon, was formed in September 2016 to formulate current evidence-based clinical practice guidelines for JTF. A systematic literature search was carried out using PubMed, the MEDLINE, and Cochrane Database of Systematic Reviews from 1982 to November 2018 applying the terms “jejunal, postpyloric, transpyloric, jejunostomy, feeding, nutrition, food.” References in these documents were also searched to ensure acquisition of relevant source data. Grading of Recommendations, Assessment, Development, and Evaluation was applied to evaluate the outcomes. Levels of evidence for each statement were based on the grading of the literature. Using the GRADE system, the quality of evidence was graded as follows (13–18).

1. High: Further research is unlikely to change our confidence in the estimate of effect.

2. Moderate: Further research is likely to have impact on our confidence in the estimate of effect and may change the estimate.
3. Low: Further research is likely to have an impact on our confidence in the estimate of effect and likely to change the estimate.
4. Very low: Any estimate of effect is uncertain.

The strength of recommendations (SoRs) was defined as follows:

Strong: when the desirable effects of an intervention clearly outweigh the undesirable effects, or they clearly do not. It should be noted that the expert group can make strong recommendations based on lesser evidence when high-quality evidence is impossible to obtain and the anticipated benefits strongly outweigh the harms. Strong recommendations are formulated as “the working group recommends (...).”

Weak: when the trade-offs are less certain (either because of the low quality of evidence or because the evidence suggests that desirable and undesirable effects are closely balanced). Weak recommendations are formulated as “the working group suggests(...).”

The ESPGHAN WG anonymously voted on each recommendation. A 9-point scale was used (1 strongly disagree to 9 fully agree), and votes are reported for each recommendation. It was decided in advance that consensus was reached if >75% of the WG members voted 6, 7, 8, or 9. Consensus was reached for all questions. In the absence of evidence from randomized controlled trials (RCTs), the majority of recommendations reflect the expert opinion of the authors. The final draft of the clinical guideline was sent to all the committee members for approval in December 2018, and then critically reviewed by a multidisciplinary panel of the GI and Nutrition committees and members of the council of ESPGHAN.

Q1: What are the indications for jejunal tube feeding?

1. The ESPGHAN expert group recommends jejunal feeding as the route of choice for providing enteral nutrition (EN) in children with failure of oral and intra-gastric feeds or gastric outlet obstruction.

Level of evidence (LoE): very low

SoR: strong

Vote: 9,9,9,9,9,8,9,7,8,9,9,8,9,9,9 (100% agreement)

2. The ESPGHAN expert group recommends that transpyloric tube feeding be considered to provide EN when gastric feeding fails in critically ill children.

LoE: moderate

SoR: weak

Vote: 9,9,9,9,9,9,9,8,9,9,9,9,9,9,9 (100% agreement)

|||Pediatric Gastroenterology, Hepatology and Nutrition, La Fe University Hospital, Valencia, Spain, the ¶¶Pediatric Gastroenterology, Grangelettes Clinic, Geneva, Switzerland, the ##The Medical College of Wisconsin, Milwaukee, WI, the ***Pediatric Gastroenterology Unit, Hadassah – Hebrew University Medical Center, Jerusalem, Israel, and the †††Neurogastroenterology and Motility Unit, Department of Pediatric Gastroenterology, Great Ormond Street Hospital NHS Foundation Trust, London, UK.

Address correspondence and reprint requests to Ilse J. Broekaert, MD, University Children's Hospital Cologne, Cologne, Germany and Nikhil

Thapar, MD, Neurogastroenterology and Motility Unit, Department of Pediatric Gastroenterology, Great Ormond Street Hospital NHS Foundation Trust, London, United Kingdom (e-mail: ibroekaert@gmail.com, n.thapar@ucl.ac.uk).

Ilse J. Broekaert, and Nikhil Thapar are shared first co-authorship. The authors report no conflicts of interest.

Copyright © 2019 by European Society for Pediatric Gastroenterology, Hepatology, and Nutrition and North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition

DOI: 10.1097/MPG.0000000000002379

3. The ESPGHAN expert group recommends that a trial of JTF be considered in children with paediatric intestinal pseudo-obstruction who fail gastrostomy feeding.
LoE: moderate

SoR: strong

Vote: 9,8,7,9,9,8,9,7,8,9,9,9,9,9 (100% agreement)

Practical note

Especially if there is some evidence of propagative peristalsis JTF should be considered.

4. The ESPGHAN expert group recommends to consider JTF as an alternative to fundoplication and gastrostomy tube feeding in children with severe gastroesophageal reflux with risk of aspiration (eg, neurological disability).
LoE: moderate

SoR: strong

Vote: 9,8,8,9,8,9,9,8,9,8,9,9,9,8 (100% agreement)

Practical note

Gastroesophageal reflux or risk for gastroesophageal reflux worsening is not a contraindication for JTF unless JTF worsens gastroesophageal reflux.

5. The ESPGHAN expert group recommends the use of JTF in children with acute pancreatitis only in cases in which oral or gastric feeding is not tolerated.
LoE: moderate

SoR: strong

Vote: 9,8,9,9,8,9,7,8,9,9,9,9,9,8 (100% agreement)

In general, the choice of the route of enteral feeding depends on several major criteria, that is, the duration of EN support, the integrity and functioning of the upper GI tract, and the risk of aspiration. In 2010, the ESPGHAN Committee on Nutrition recommended that postpyloric feeding is indicated only in clinical conditions in which gastroparesis/dysmotility, aspiration, gastric outlet obstruction, or previous gastric surgery precludes gastric feeding or when early postoperative feeding after major abdominal surgery is planned (6). The evidence to support these recommendations is not based on controlled studies.

Since 2010 some studies and guidelines were published concerning indications of jejunal and postpyloric feeding in different clinical situations.

GASTRIC DYSMOTILITY: CRITICALLY ILL CHILDREN, PRETERM INFANTS, CHRONIC INTESTINAL PSEUDO-OBSTRUCTION, GASTROPARESIS, AND SHORT BOWEL SYNDROME

Critically Ill Children

In accordance with the 2010 ESPGHAN recommendations, the American Society for Parenteral and Enteral Nutrition (ASPEN)/Selection and Care of Central Venous Access (SSCM) and European Society for Clinical Nutrition and Metabolism (ESPEN) guidelines advise against routine use of postpyloric feeding in the adult critically ill patient unless the patient has a high risk for aspiration or gastric feeding intolerance (8). In critically ill children, the ASPEN guideline (2009) states that postpyloric feeding should be considered in patients at high risk of aspiration or in whom gastric feeding fails (9). Both international bodies recognized that there is limited research data available.

In a systematic review and meta-analysis looking at gastric versus postpyloric feeding in critically ill adults moderate- to low-quality evidence was found showing a lower rate of ventilator-associated pneumonia with postpyloric feeding (19,20) and low-quality evidence suggesting an increase in the amount of nutrition delivered to these participants (19). Because no differences were found between gastric and postpyloric feeding for objective outcome measures such as mortality, duration of mechanical ventilation and complications, the finding of decreased ventilator associated pneumonia in postpyloric feeding may not be real (20).

In 2014, an international survey among 31 paediatric intensive care units evaluating institutional nutrition practices showed that 13.2% of patients were postpylorically fed, but only 9 units had detailed EN algorithms (21). All recommended the use of postpyloric feeding where gastric feeding has failed and/or where concerns about pulmonary aspiration exist (21).

A meta-analysis from 2013 (22) comparing the use of postpyloric versus gastric feeding in adults and children in the ICU, including 17 trials (1 paediatric RCT with 30 patients) (23), showed that postpyloric feeding, overall, delivered significantly more nutrition than gastric feeding, with a weighted mean difference of 12%. The meta-analysis failed to demonstrate any benefits of postpyloric feeding with regards to new-onset pneumonia, mortality, and aspiration (22).

In a Cochrane review in 2016 regarding nutritional support in critically ill children no studies addressed JTF.

In conclusion, JTF can be a good option for providing EN when gastric feeding fails in critically ill patients. There is conflicting evidence about the prevention of complications and studies in critically ill children are lacking.

Preterm Neonates

A Cochrane review on preterm infants updated in 2013 (24) with a total of 9 RCTs (359 premature infants, studies from 1975 to 1992) failed to show beneficial effect of transpyloric feeding on feed tolerance or in-hospital growth. This is discussed in detail in question 4.

Paediatric (Chronic) Intestinal Pseudo-obstruction

In a prospective study of JTF in children with chronic intestinal pseudo-obstruction 18 children dependent on parenteral nutrition (PN) and failing gastric feeding were initiated on elemental feeding via surgical jejunostomy after performing antroduodenal manometry (25). Follow-up showed that 12 of these children (9 with and 3 without migrating motor complexes on manometry) tolerated JTF well and PN could be stopped. Although not specifically addressed, an ESPGHAN guideline from 2018 recommended that strategies such as JTF could be considered in patients with intestinal pseudo-obstruction (7).

Gastroparesis

Gastroparesis in children is most often idiopathic with other causes including, post viral and drug-related issues and occurring in association with comorbidities. It is characterized by delayed gastric emptying of solids and/or fluids without evidence of a mechanical gastric outlet obstruction. JTF may be indicated in the management of gastroparesis in cases when medical therapies fail and when nutritional intake is inadequate (26–28).

Short Bowel Syndrome

Jejunal feeding can be considered in children with short bowel syndrome in case of severe GERD, or severe gastric or upper intestinal dysmotility when oral or gastric enteral feeding fails. The limiting factor, however, will be the fact that with jejunal feeding a substantial part of the small bowel will be bypassed, thereby impairing the process of intestinal adaptation and further decreasing the absorptive capacity that is already limited in short bowel syndrome. Furthermore, the presence of a jejunal feeding tube may increase the risk of intestinal contamination with a change of the gut microbiome and subsequent small intestinal bacterial overgrowth (SIBO). However, if JTF is the only option using the enteral route, it can be considered to induce intestinal adaptation (29,30).

SEVERE GASTROESOPHAGEAL REFLUX WITH RISK OF ASPIRATION (EG, NEUROLOGICAL DISABILITY)

GERD and swallowing problems are common in children with neurological impairment (NI) and predispose to aspiration pneumonia, which is the most common cause of death in these children. They often require fundoplication and gastrostomy tube placement. Various studies have retrospectively looked at JTF as an alternative option for treatment of GERD, but RCTs and prospective studies are lacking.

The 2017 ESPGHAN guideline suggests the use of JTF where there is a risk of aspiration due to GERD (5).

A systematic review and meta-analysis specifically in children with NI, included retrospective studies of GJT versus fundoplication with gastrostomy in the management of severe GERD (31). Of these, 3 studies reporting 556 children (fundoplication with gastrostomy [$n=431$] and GJT [$n=125$]), showed no differences in rates of pneumonia (17% vs 19%) or mortality (13% vs 14%) (32–34). Furthermore, no statistically significant differences were found between the occurrence of major complications (fundoplication with gastrostomy (29%) compared to GJT (12%) (risk ratio = 1.70, 95% confidence interval [CI] 0.85–3.41, $P=0.14$)) and minor complications (GJT [70%] vs fundoplication with gastrostomy [45%], risk ratio = 0.38, 95% CI 0.05–3.07, $P=0.36$). No studies reported on quality of life using validated measures. The authors concluded that because of very low quality of evidence, large comparative studies are needed to find out which approach is associated with the best quality-of-life outcomes.

A number of more recent retrospective studies looking at short- and long-term outcomes of GJT feeding in children with NI and GERD have suggested that although major complications are comparable to fundoplication, GJT feeding is associated with reasonable amounts of morbidity (1–3,35).

In a study on pathophysiology by Rosen et al (35) in which transpyloric feeding as an alternative treatment of gastroesophageal reflux (GER) was evaluated, multichannel intraluminal impedance tracings showed that reflux events, although significantly less than previously reported in patients with significant GERD, were still present especially during feeding periods. Furthermore, patients continued to have the same amount of aspiration events and reflux-related hospitalizations after start of transpyloric feeding.

A large retrospective study in children with NI and GER requiring gastrostomy tube feeding who either underwent initial GJT placement ($n=163$) or fundoplication ($n=1178$) showed that first-year postprocedure reflux-related hospitalization rates, and odds of death were similar in both groups, whereas failure to thrive, repeat of initial intervention, and crossover intervention were more common in the GJT group (36). It was concluded that either intervention could reduce future aspiration risk; the choice can

reflect nonreflux-related complication risks, caregiver preference, and clinician recommendation.

Egnell et al (2) reported retrospectively on the clinical outcome and safety of surgically placed jejunostomies in 33 children (of which 17 with NI). They concluded that these types of tubes could be effective and safe in selected children with GERD, feeding difficulties, or recurrent pneumonia.

GASTRIC OUTLET OBSTRUCTION

The use of JTF in case of upper GI obstruction has been studied mostly in adults with gastric or pancreatic cancer and benign pancreatic diseases. Few studies have been performed in children and prospective RCTs are lacking. A retrospective study in 120 children who had undergone surgery for duodenal and jejunal congenital obstructions (ie, duodenal atresia, annular pancreas, jejunal atresia) showed that children in the early EN group with feeding through a nasojejunal tube (NJT) had a better outcome compared to children in the control group on PN (37). The JTF group experienced a shorter time to tolerate oral feeding and a lower incidence of cholestasis and had a shorter postoperative hospital stay. Another retrospective study from the same research group showed that feeding through an NJT could safely be provided in neonates after partial gastrectomy ($n=46$) because of gastric perforation and led to fewer complications than total PN (38).

ACUTE PANCREATITIS

The use of JTF in patients with severe acute pancreatitis is mostly performed in adult patients. In children, no RCTs are available and no guidelines have been published regarding type or route of nutritional support in acute pancreatitis.

The North American Society for Pediatric Gastroenterology, Hepatology and Nutrition (NASPGHAN) recently published a clinical report about the management of acute pancreatitis in the paediatric population with conclusions mostly based on adult literature. Recommendations include early EN as tolerated, whether through oral, gastric, or jejunal route (39).

Q2: What alternatives can be tried before finally considering jejunal feeding?

6. The ESPGHAN expert group recommends a trial of continuous gastric feeding with a hydrolysed or elemental formula before postpyloric feeding is started.

LoE: high

SoR: strong

Vote: 9,7,9,9,9,5,9,6,8,7,7,9,7,7 (87% agreement)

7. The ESPGHAN expert group recommends to consider trialling at least 1 prokinetic drug to promote oral or gastric feeding before instituting jejunal feeding as they are widely used, but there is no published data.

LoE: moderate

SoR: strong

Vote: 9,7,9,7,8,8,9,6,8,6,7,8,9,3,8 (80% agreement)

Meyer et al carried out a systematic review on the impact of feed protein type and degree of hydrolysis on gastric emptying in children. Although this was limited by considerable variability between the studies, a number of studies reported better emptying by hydrolysed compared to whole protein (40).

After other aetiologies are excluded and before a drug trial, gastric feeding with a hydrolysed or elemental formula should be performed.

Although a number of pharmacological therapies are used to improve gastric emptying and feeding tolerance, published paediatric literature is scarce and only a few controlled trials exist, whereas none of them has addressed the issue of drug introduction *before* jejunal feeding. Nearly all the published studies were performed on premature infants with the goal of advancing EN as quickly as possible. Most studies were retrospective. A recent guideline on feeding premature infants did not even discuss drug therapy (41). Available agents include erythromycin, a macrolide antibiotic and nonpeptide motilin agonist, as well as metoclopramide and domperidone, both dopamine D2 receptor antagonists.

ERYTHROMYCIN

Intragastric or intravenous administration of a low dose (3–5 mg/kg/dose 3–4 times daily) of erythromycin induces a migrating motor complex. Although a number of studies in premature infants suggests improved feeding tolerance in subjects given erythromycin compared with control infants a 2008 Cochrane review concluded that there is insufficient evidence to recommend the use of erythromycin in low or high doses for preterm infants with or at risk of feeding intolerance (42).

A recent retrospective multicentre review of 1095 infants treated with erythromycin and 19001 infants treated with metoclopramide stated that “*the safety and efficacy of erythromycin in infants is incompletely characterized*” (43) but that there were fewer adverse events with erythromycin than with metoclopramide.

Although frequently quoted there is no published data suggesting that the use of erythromycin benefits a paediatric patient intolerant of gastric feeding except for small single-centre studies in premature infants (44).

In adult studies erythromycin may be more effective than metoclopramide for gastroparesis (28) but this may not be true in children (27). Tachyphylaxis to the prokinetic effect of erythromycin develops within 4 weeks.

METOCLOPRAMIDE

An RCT performed by Hyman et al (45) found that metoclopramide is not efficacious in premature and neonatal populations whose primary cause of gastroparesis is prematurity. Tube feedings that contained 0.2 mg/kg metoclopramide had no effect on promoting gastric motility in low-birth-weight neonates but may be helpful in reducing emesis due to its actions on the chemoreceptor trigger zone. The usefulness of metoclopramide in neonates may be due to the centrally acting antiemetic properties and not the prokinetic effect seen through binding of the dopamine D2 receptor in the peripheral nervous system. Another study showed that 80% of paediatric patients with gastroparesis failed to respond to metoclopramide therapy (46).

Because of the risk of tardive dyskinesia, the US Food and Drug Administration issued a black box warning for metoclopramide in 2009. In 2013 the European Medicines Agency restricted usage to children over 1 year of age and for a duration no longer than 5 days. The suggested dose of metoclopramide is 0.4 to 0.8 mg · kg⁻¹ · day⁻¹ 30 minutes before feeding.

DOMPERIDONE

Domperidone is available in many European countries and in Canada, but not in the United States. In some countries it is available over the counter. It is considered less safe than erythromycin. In adults it is more effective than metoclopramide for gastroparesis (47). The only paediatric studies relate to its use for GER and as an antiemetic. In 2014 the European Medicines

Agency restricted use to the treatment of nausea and vomiting. The suggested dose of domperidone is 0.1 to 0.3 mg/kg per dose 2 to 4 times daily 30 minutes before feeding.

Q3: What investigations should be carried out before jejunal tube feeding placement?

8. The ESPGHAN expert group recommends to consider performing a contrast meal and follow through study of the small intestine in all patients to ensure patency of the intestinal lumen and exclude a mechanical obstruction before jejunal feeding tube placement.

LoE: very low

SoR: strong

Vote: 9,9,9,7,9,9,5,8,7,8,8,9,9,7 (93% agreement)

9. The ESPGHAN expert group recommends to consider an upper GI endoscopy in all patients before concomitant to JTF placement.

LoE: moderate

SoR: strong

Vote: 9,8,9,8,9,8,8,9,8,9,8,9,9,7,9 (100% agreement)

Clearly in the lead up to JT placement in children there should be careful consideration of the rationale for using this route of feeding including the reasons for the failure of oral or gastric feeding and any contraindications for postpyloric feeding. It is these considerations that essentially underlie what investigations should be carried out before postpyloric tube placement. Several excellent guideline papers are available on the use of condition-specific investigations.

Of particular importance are investigations, largely imaging that provide information about the patency of the GI tract and presence of any mechanical problems and about previous GI tract surgery and risk of intestinal perforation (Table 1). In patients suspected of gastroparesis for example diagnostic evaluation may include an upper endoscopy to rule out mechanical causes, followed by a gastric-emptying scintigraphy for diagnosis. Other diagnostic alternatives that have been used include wireless capsule motility, antroduodenal manometry, and breath testing (48). Apart from these there is virtually no evidence from the available literature for the routine application of a battery of investigations before the placement of a postpyloric (jejunal) feeding tube.

In theory, if not applied previously, a contrast follow through study of the small intestine should be carried out to ensure patency of the intestinal lumen and exclude a mechanical obstruction or issue that may increase the risk of intestinal perforation or impaired viability (eg, intestinal pneumatosis). There is some evidence to suggest other investigations may add diagnostic value in particular groups of patients.

Van Haren et al (49) carried out a retrospective observational case study on adult intensive care patients who underwent postpyloric feeding tube insertion under endoscopic guidance. They found significant endoscopic findings in almost 50% of the patients in whom endoscopic reports were available and suggested that endoscopic placement of postpyloric feeding tubes resulted in the identification of a significant number of patients with previously undiagnosed upper GI tract abnormalities (49). Others have similarly shown that diagnostic upper endoscopy performed

TABLE 1. Investigations recommended before placement of postpyloric feeding tube

	Consider routinely in all patients before placement of postpyloric feeding tube	Consider on case-by-case basis to rule out underlying disorders (in parentheses) that may limit prepyloric feeding
Investigation	Upper GI endoscopy (<i>upper GI abnormalities eg, esophagitis, ulceration, lesions causing obstruction not picked up on contrast studies</i>) Contrast meal and follow-through (<i>mechanical obstruction</i>)	pH/impedance studies (<i>gastro-oesophageal reflux disease</i>) Gastric emptying (nuclear medicine/scintigraphic) studies (<i>gastroparesis</i>) Antroduodenal manometry (<i>paediatric intestinal pseudo-obstruction</i>)

GI = gastrointestinal.

concomitantly with placement of the JT often reveals findings of clinical importance (50,51).

Q4: What are the absolute and relative contraindications against jejunal tube feeding?

10. The ESPGHAN expert group recommends to avoid JTF in the presence of the following conditions: paralytic or mechanical ileus, intestinal obstruction, intestinal perforation, peritonitis, and necrotizing enterocolitis (NEC; Table 2: absolute contraindications).

LoE: very low

SoR: strong

Vote: 9,9,7,9,9,9,8,9,9,9,9,9 (100% agreement)

11. The ESPGHAN expert group recommends caution when considering JTF in patients with relative contraindications to transpyloric feeding: preterm infants, intestinal dysmotility, toxic megacolon, GI bleeding, high-output enteric fistula, intractable diarrhoea, immunocompromised children (Table 2: relative contraindications).

LoE: moderate

SoR: strong

Vote: 9,9,9,9,9,9,8,9,9,8,9,8 (100% agreement)

12. The ESPGHAN expert group recommends not to use JTF in preterm infants (<37 weeks' gestation).

LoE: moderate

SoR: strong

Vote: 9,8,8,9,8,9,5,8,9,8,9,9,9 (93% agreement)

A systematic review of transpyloric versus gastric tube feeding for preterm infants (<37 weeks gestation) came to the conclusion that because of the lack of evidence of any benefit, and an increased risk for GI disturbance and possibly of death the transpyloric route should not be routinely used for preterm infants who require enteral tube feeding (52). Especially preterm infants with intrauterine growth restriction are at a higher risk for adverse events. Most of the studies recruited very-low-birth-weight infants (birth weight <1500 g), although in the majority only infants grown appropriately for gestational age were included (52).

Two additional systematic reviews and the most recent Cochrane review conclude that there is no evidence of any benefit for transpyloric feeding in preterm infants compared to gastric feeding (24,53,54). In addition, a higher risk for GI complications (relative risk 1.45, 95% CI 1.05, 2.09), and a higher mortality rate

(relative risk 2.46, 95% CI 1.36, 4.46) before discharge from the hospital was observed in preterm infants fed transpylorically. However, because of allocation bias in the included trials—sicker and less mature infants were allocated to JTF—the authors of the systematic review advise that these findings should be interpreted with caution (24). Nevertheless, the ESPGHAN committee on nutrition recommends to avoid postpyloric feeding in preterm infants (6). It has been suggested that early transpyloric feeding in preterm infants may prevent bronchopulmonary dysplasia, but this must be further tested in studies before it can be recommended (55).

Other conditions considered relative contraindications include intestinal dysmotility, toxic megacolon, peritonitis, GI bleeding, high-output enteric fistula, and intractable diarrhoea (6). These are not deemed absolute contraindications as minimal quantities of nutrients in the GI tract—so-called trophic feeding—have well recognized benefits by promoting intestinal perfusion, initiating release of enteral hormones and improving gut barrier function (56,57). Severe vomiting may compromise the benefits of JTF or impair viability, also through tube displacement (6). JTF is not a contraindication in patients with evidence of gastrooesophageal reflux (GOR) or risk for GOR worsening, for example, in children with severe NI. Placing the tip of the JT beyond the ligament of Treitz prevents duodenogastric reflux and GOR and this is suitable for children who are not acceptable candidates for antireflux surgery or in whom fundoplication has failed (10). This is discussed in detail in Question 1.

If long-term enteral feeding is required the high frequency of complications and the need for frequent tube replacement due to obstruction or displacement could be a limitation to JTF (10–12). However, other authors have reported enteral feeding through surgically placed jejunostomy tubes to be relatively safe even for long periods up to 12 years (2).

As placement of nasoduodenal or NJTs may be difficult percutaneous endoscopic gastrostomy (PEG) or percutaneous endoscopic enterostomy currently are the preferred routes of placement

TABLE 2. Absolute and relative contraindications to jejunal tube feeding

Absolute contraindications	Relative contraindications
Paralytic ileus	Preterm infants
Mechanical ileus	Intestinal dysmotility
Intestinal obstruction	Toxic megacolon
Intestinal perforation	Peritonitis
Necrotizing enterocolitis	Gastrointestinal bleeding
	High-output enteric fistula
	Intractable diarrhoea
	Immunocompromised children

especially for long term EN (6). Here, contraindications for PEG and percutaneous endoscopic enterostomy need to be considered (6).

Other conditions may be considered as limitations for JTF because of higher rates of significant complications. In immunocompromised children or in case of an impaired gastric acid barrier, there may be a higher risk for sepsis from bacterial contamination of feeds which is relative common both at home and during hospitalization (58). As patients being JT fed have a 9 times higher risk for developing *Clostridium difficile*-associated diarrhoea as compared to matched controls, decision of JTF in immunocompromised individuals needs to be carefully considered (59).

JTF can, however, be safely used in children on chronic ventilation or during weaning of mechanical ventilation as no higher risk for aspiration or mortality has been noticed (60,61).

The presence of a ventriculoperitoneal shunt or a peritoneal dialysis catheter is not a contraindication to JTF as a PEG does not result in a higher incidence of shunt infections or mortality (62); however, it has been suggested that PEG insertion should be deferred at least 1 week after ventriculoperitoneal shunt insertion (62).

Q5: What are the techniques available for placement of a jejunal feeding tube?

The route of placement and type of device used for jejunal feeding should depend on the expected duration of jejunal feeding, namely NJT for predictably <1 month and per endoscopic or surgical gastrojejunostomy or jejunostomy for more prolonged use; the availability of experience and collaboration locally (ie, interventional radiology, surgery, and endoscopy); the presence of a pre-existing gastrostomy; and the need of gastric decompression.

Nasajejunal Tube

13. The ESPGHAN expert group recommends that the radiological placement of an NJT should follow established protocols and training of clinical staff to reduce radiation exposure of patients.

LoE: low

SoR: strong

Vote: 9,9,9,9,9,9,8,9,8,9,9,9,9 (100% agreement)

14. The ESPGHAN expert group suggests not to use prokinetic agents during nasajejunal feeding tube placement.

LoE: moderate

SoR: weak

Vote: 9,7,8,9,9,8,8,5,8,9,8,9,7,8 (93% agreement)

Nasoenteric tubes are a good choice for short-term feeding but have many drawbacks for long-term management (recoil into the stomach, clogging, nasal pressure sores, and accidental removal). There are several kinds of nasoenteric tubes made from various materials (eg, polyurethane and silicone), which have different diameters (3.5–12 French), with and without guidewires, and with and without weight at their tips. Nasogastric tubes made of polyvinyl chloride are relatively stiff and therefore more irritating long-term, and are used primarily for GI decompression and should not be used for prolonged enteral feeding. Usually, a nasoenteric tube is inserted with a guidewire, previously flushed with saline solution for easier postinsertion wire removal, and a weighted tip is inserted into the stomach using the usual technique for nasogastric tube insertion. The child is placed in right lateral decubitus and the tube is pushed through

the pylorus. The guidewire should be removed at the end of the procedure. Self-advancing jejunal feeding tubes have been reported to be used effectively to establish early EN in critically ill children (63). The use of a noninvasive electromagnetic device to place transpyloric feeding tubes has been suggested to be effective in children (64), although in another study it significantly increased the time of placement (65). Several studies showed that the insufflation of 10 mL/kg air in the stomach significantly improves the rate of success without increasing risks (66,67). A recent meta-analysis of all the RCTs both in adults and children concluded that gastric air insufflation seems to be efficient (without reaching significance), whereas clinicians should no longer use prokinetic agents in paediatric patients or patients without impaired motility (66,67). Bedside placement of a postpyloric tube can be safe and effective in infants including preterm and reduce infants' exposure to radiation in comparison to interventional radiology placement (68). Nasoenteric tubes may also be placed with the aid of fluoroscopy or endoscopy. Fluoroscopic techniques of nasoenteric tube placement require skilled radiological support and cause exposure to radiation. Protocols and training can reduce radiation exposure of patients and staff (69). The NJT can be placed endoscopically, either using a guidewire introduced through the working channel of the gastroscope or the drag technique in which a suture is tied to the end of a feeding tube and dragged with the endoscope snare or forceps from the stomach to the duodenum. This procedure is less successful because the feeding tube frequently moves back into the stomach when the endoscope is removed unless the tip of the tube is clipped in the duodenum (this is limited to older children due to the opening size of the clip). Irrespective of the technique used for NJT placement, proper position of the nasoenteric feeding tube must be verified radiographically before feeding is initiated.

pH-guided Jejunal Tube Placement

15. The ESPGHAN expert group recommends to use pH-guided jejunal feeding tube placement whenever possible as a safe, easy, and cost-effective bedside method.

LoE: low

SoR: strong

Vote: 9,8,9,9,8,8,7,8,9,8,8,9,9,8 (100% agreement)

pH in the upper GI tract typically varies according to the anatomical segment (oesophagus: pH 5–7, stomach: pH 1–3, duodenal bulb: pH 3–4 and small intestine: pH 7–8) (70). Therefore, pH-guided JT placement is a safe, easy, and cost-effective bedside alternative to fluoroscopic, endoscopic, or surgical placement in critically ill infants and small children (70,71). This method can be easily taught to house staff or other health care personnel (70). As the pH-assisted technique offers immediate feedback on correct positioning enteral feeding can be initiated promptly (71). Displacements of jejunal feeding tubes can be easily checked with a pH monitor, and therefore, aspiration of jejunal secretions to check pH with paper is not needed (71). Radiological placement control should only be applied in case of borderline pH values or in patients treated with proton pump inhibitors.

RADIOLOGY

Radiological methods can help in placing NJTs and are needed to confirm proper tube position. Jejunal feeding tubes

can also be placed under radiological guidance via a previous gastrostomy site (72) or by direct jejunal puncture (73). A retrospective review comparing surgical jejunostomy against image-guided GJT placement through a pre-existing gastrostomy orifice concluded that image-guided GJT placement needed more frequent tube replacement (4.6/year vs 1.5/year) ultimately leading to surgical jejunostomy conversion in 50% of the cases (74).

ENDOSCOPY

A jejunostomy may be inserted with endoscopic assistance indirectly *via* a previously placed or a *de novo* gastrostomy (percutaneous endoscopic gastrojejunostomy [PEG-J]) or directly without gastrostomy placement (percutaneous endoscopic jejunostomy [PEJ]).

For PEG-J placement, a feeding tube long enough to pass beyond the pylorus is inserted through an existing gastrostomy. The tip of the feeding tube is then grasped with the biopsy forceps of the endoscope and the tube is pushed into the duodenum as far as possible. Extra tubing length is left within the stomach to allow peristalsis to pull the tip of the feeding tube past the ligament of Treitz. Although this procedure is simple, its major disadvantage is the tendency of the feeding tube to recoil into the stomach during the withdrawal of the gastroscopy; a clip can limit this risk fixing the external part of the tube to the duodenum/jejunum (this is limited to older children due to the opening size of the clip). In addition, the feeding tube tends to dislodge from the outer gastrostomy. An alternative is to introduce a neonatoscope (diameter 5.3 mm) through the gastrostomy, pass the pylorus, and go as far as possible beyond the Treitz angle. Then a guide wire is introduced through the operating channel of the endoscope, the scope is removed and the GJT is passed over the guidewire (12). One advantage of this technique is that it minimizes the need for sedation because it causes minor discomfort such as hiccups, pain around the stoma site, and abdominal distension from air insufflation. These can be overcome in most patients by providing play therapy and the presence of the parents during the procedure (75). One-step GJT insertion through a *de novo* gastrostomy is a recent technique using the push technique. The procedure is basically the same as the 1-step percutaneous endoscopy button placement (76) where a neonatoscope is introduced through the 16-French introducer and passed into the jejunum via the pylorus and a GJT is placed over the guidewire as described above (77).

If there is no pre-existing gastrostomy or if a gastrostomy (for exsufflation or administration of medication) is not needed, direct PEJ can be performed using a gastroscope or colonoscope placed into the proximal jejunum. The most common techniques include the insertion of a needle into the jejunal lumen at the site of the maximal transillumination and/or a finger indentation marking of the jejunal loop that is closest to the abdominal wall. The needle should be snared tightly, fixing the small bowel against the abdominal wall. The plastic sheath with stylet should then be inserted adjacent to the needle and snared by a wire loop that has been removed from the needle. An insertion wire is then passed through the plastic sheath and grasped with a snare or a grasp forceps. The rest of the procedure is similar to the PEG's pull technique: the gastroscope with a wire is pulled out through the duodenum, stomach, oesophagus, and mouth. The insertion wire is then secured to the loop at the end of the feeding tube with an internal jejunal bolster and the assembly is pulled through the mouth all the way to the jejunum. The tube is pulled through an incision in the abdominal wall, sufficiently tight to compress the jejunal wall against the anterior abdominal wall. Intrajejunal tube placement is then verified by a second endoscopy. Finally, a skin disk is secured to the outside portion of the feeding tube to ensure the creation of a tract between the skin and jejunal lumen. It is important to avoid excess tension when approximating the jejunum to the abdominal wall and to prevent pressure sores of the skin or the

jejunal mucosa. Experience in children with this technique remains limited (78). Recently a laparoscopic-assisted PEJ technique has been reported in 16 children aged 2 to 17 years. All procedures were successful and the technique was safe because it provides sufficient visualization of the bowel loops intra-abdominally (79).

Surgery

16. The ESPGHAN expert group recommends, where long-term (gastro-) jejunal feeding is expected, to use strategies such as Roux-en-Y jejunostomy, Omega jejunostomy, and retubularization instead of direct surgical tube insertion.

LoE: low

SoR: strong

Vote: 9,9,7,9,9,9,9,8,8,9,8,9,9,9 (100% agreement)

Many different surgical techniques have been described for JT insertion. Open or laparoscopic surgery techniques are available and there are no data that demonstrate superiority in effectiveness and safety of any strategy, and the choice depends on the surgeon's experience and his preferences.

Direct surgical catheter jejunostomy placement is a well-known and standardized procedure. High surgical complication rates (40%) have, however, been reported in a large series (2).

Laparoscopic side insertion of a small calibre tube (6–9 Fr) or Foley catheter into the proximal jejunal loop is a straight-forward technique (80). A subserous tube conduit prevents the risk of peristomal skin damage due to leakage and tube dislodgement.

Laparoscopic insertion of a GJT has been described in a large group of infants <10 kg with cardiac disease (81). There are no clear advantages of this strategy except for the reduction of gastric and bowel distension during the procedure.

In case of long-term JTF, surgical strategies such as Roux-en-Y jejunostomy (82), Omega jejunostomy (83) or retubularization (84) facilitate insertion of the tube into a modified jejunal tract improving management by the caregivers and also reduce peristomal leakage and skin damage.

Q6: Which complications are related to JTF and how should they be minimized and/or managed?

The development of procedural protocols with regular quality controls and audits, and monitoring by a dedicated nutrition support team warrants to minimize complications. Although GJTs are a useful temporizing method to provide enteral access in children, their high rate of mechanical failure limits their long-term use.

17. The ESPGHAN expert group recommends that the tip of the jejunal feeding tube be placed beyond the ligament of Treitz to prevent retrograde dislodgment of the tube into the stomach.

LoE: very low

SoR: strong

Vote: 9,9,9,9,9,9,9,8,9,9,9,9,9,9 (100% agreement)

There are 3 major categories of complications following JT placement: mechanical (eg, perforation, buried bumper syndrome); GI (eg, diarrhoea); infectious (eg, aspiration pneumonia, tube site infection) (Table 3).

TABLE 3. Late complications of postpyloric feeding (excluding events related to jejunal tube placement): possible causes, prevention, and treatment

Mechanical complications	Possible causes	Prevention and treatment
NJT/GJT obstruction/ clogging, knotting	Thick enteral feeds Medications Bulking agents (i.e. resins) Small lumen	Water flushing after feeding and medications Accurately dissolve medications before administration Prefer liquid drug formulations Mechanical cleaning with wires or special “declogging” brush devices Tube substitution Continuous infusion
NJT/GJT displacement (retrograde dislodgment in the stomach)	Initial positioning of the tube tip before the ligament of Treitz Altered gastrojejunal motility with no regular peristalsis	Tube tip beyond the ligament of Treitz Endoscopically placed clips Tube replacement (“beneath the scope” or “over the wire technique”)
Accidental NJT/GJT tube removal	Inadequate fixing Excessive traction of the tube during feeding Patient’s poor compliance	Appropriate fixing, specific fixing devices Avoid traction during feeding Patient/ caregiver education Contention
Breakage, leakage, wear of the NJT/GJT tube; rupture of the GJT balloon	Excessive wear Inadequate manipulation	Tube substitution Patient/caregiver education
Peristomal leakage (and subsequent erosion, ulceration and necrosis of skin and mucosa)	Infection/ bleeding at the GJT insertion site Gastric hypersecretion Excessive torsion of the tube Excessive cleansing with hydrogen peroxide Host factors for poor wound healing Inadequate size of the device Inadequate stabilization by the external bolster	Reduction of risk factors (ie, antisecretory therapy with PPIs) Barrier creams containing zinc and skin protectants Placement of a smaller diameter tube Apply continuous low pressure suction (ie, Replegle tube)
Buried bumper syndrome	Excessive traction between the internal bumper and the stomach wall	Appropriate size of the device (length according to abdominal wall thickness and weight gain)
Intestinal perforation, intussusception, intestinal obstruction	Young age - Comorbidities (ie, shock, heart disease)	
Infectious complications	Possible causes	Prevention and treatment
Infection at the GJT insertion site Peritonitis	Improper wound dressing (ie, occluding) Excessive traction between the internal bumper and the stomach wall Host factors (ie, immunosuppression)	Regular and appropriate skin and stomal care (ie, antimicrobial wound dressings) Proper size of the device (length according to abdominal wall thickness and weight gain) Topical or systemic antibiotics
Infectious diarrhoea	Inadequate manipulation and storage of feeding formula Host factors (ie, immunosuppression)	Hygienic manipulation and storage of feeding formula
NEC	Host factors (ie, prematurity, shock) Vasoactive drugs	Surgery and/or medical treatment
Jejunoileitis	Local vascular compromise Bacterial overgrowth	Surgery and/or medical treatment
Nasopharyngeal and ear infections with NJT	Partial upper airway obstruction by NJT	Substitution of NJT with GJT in case of prolonged postpyloric feeding
Gastrointestinal complications	Possible causes	Prevention and Treatment
Diarrhoea	Too rapid infusion rate Too cold feed temperature Hyperosmolar feedings Fat malabsorption Milk-protein intolerance Lactose intolerance Drugs	Reduce/ control infusion rate Increase to room temperature Use isotonic feeding solution, initially dilute hyperosmolar feeding solutions Low-fat or MCT-containing diet Protein hydrolysate/elemental formula Low-lactose or lactose-free diet Fibre/probiotics

Gastrointestinal complications	Possible causes	Prevention and Treatment
Persistent GERD	Underlying disease (ie, neurological impairment, oesophageal atresia, prematurity, etc)	
Respiratory complications		Possible causes
Aspiration pneumonia		NJT + supine position: combination of gravitational back-flow and presence of the tube across the gastric cardia Neurological impairment Persistent GERD
Metabolic complications		Possible causes
Refeeding syndrome		Chronic/ severe malnutrition Prolonged fasting
Overhydration		Excessive enteral + intravenous fluid intake
Electrolyte disturbances		Underlying metabolic diseases (ie, diabetes mellitus and renal/ hepatic insufficiency)
Hyper- and hypoglycaemia		Dumping syndrome: high-volume, highly refined carbohydrate in the small bowel Underlying metabolic diseases (ie, diabetes mellitus and renal/ hepatic insufficiency)
Vitamin and trace element deficiency		Pre-existing condition or inadequate intake with feeding formula, side effects of medication (eg, cholestyramine)

GERD = gastroesophageal reflux disease; GJT = gastrojejunal tube or gastrojejunostomy tube; MCT = medium-chain triglyceride; NEC = necrotizing enterocolitis; NJT = nasojejunal tube; PPI = proton pump inhibitor.

MECHANICAL AND SURGICAL COMPLICATIONS (EG, PERFORATION, INTUSSUSCEPTION, AND BURIED BUMPER)

NJT is mainly used for short-term postpyloric feeding (4–6 weeks). Its complications include foreign body sensation, obstruction, tendency to dislocate and easy voluntary removal, reflux esophagitis, aspiration, nasopharyngeal ulcers, and epistaxis (59).

Placing the distal tube tip beyond the ligament of Treitz minimizes retrograde dislodgement of the jejunal extension tube. Endoscopically placed clips may secure the tube and prevent migration.

Peristomal leakage may be reduced by adequate stabilization of the external bolster by a dressing. If persistent leakage causes peristomal skin damage, barrier creams may be helpful, as well as local antibiotics. It is essential to prevent stoma enlargement. Sometimes a smaller tube may facilitate healing around the tube, and the temporary application of continuous low-pressure suction at the insertion site (ie, negative-pressure wound therapy [VAC[®] therapy], Replogle tube).

Buried bumper syndrome may complicate the placement of a GJT, when there is excessive traction between the internal bumper and the stomach wall as, for example, in patients with important weight gain. There is a higher rate of buried bumper syndrome associated with PEG-J tubes compared to PEG tubes, possibly related to the jejunal extensions leading to difficulty in the usual maintenance regimen that all carers are taught after PEG/PEG-J insertion (85). To prevent buried bumper syndrome, it is advisable to allow some space between the external bumper of the PEG tube and the skin to minimize the risk of pressure-induced necrosis and to

mobilize and loosen the PEG from the outside at least every other day to avoid mucosal overgrowth of the inner bumper. To prevent this event, the size of the device must be reviewed periodically for weight gain and increased abdominal wall thickness (86).

Intestinal perforation may occur even much time after placement, mainly at a younger age (1) and in patients with comorbidities, that is, shock or heart disease (87,88). Intussusception has also been reported as a rare complication (73).

Peristomal infections occur more frequently shortly after first tube placement (PEG or PEJ), but may also complicate long-lasting enteral feeding. Accurate hygiene measures of the stoma and the use of antimicrobial wound dressings may help in prevention (86). Depending on clinical status, topical or systemic antibiotics may be required.

GASTROINTESTINAL COMPLICATIONS

Although GJT feeding usually improves nutritional status, its use may be associated with pulmonary aspiration, bilious aspirates, and diarrhoea (10). Diarrhoea is the most commonly reported GI side effect in patients receiving JTF. The pathogenesis of diarrhoea in enterally fed patients can be related to the enteral formula or the administration method. Prevention of diarrhoea includes the use of a closed feeding system (to limit bacterial contamination), continuous administration of feeding using a pump, and limiting the use of hyperosmolar feeds. Persistent vomiting and retching are described in almost 18% of a large series of children with GJ tube (1), but is probably more likely due to the underlying disease (severe GOR, antropyloric dysmotility, etc). The frequency of GI complications is higher in critically ill

children (89) and patients with cyanotic heart disease, which in turn increases the risk of NEC (87,88).

INFECTIOUS COMPLICATIONS (EG, ASPIRATION PNEUMONIA, TUBE SITE INFECTION)

The combination of gastric decompression via PEG and simultaneous jejunal nutrition reduces tube feeding–related aspiration in many patients. Tube site complications including granulation, infection, and leakage are frequent and benign complications. Leakage of bile acids at the level of a jejunostomy can be responsible for severe and painful skin lesions due to the caustic nature of the bile.

Q7: Immediately after placement when should feeding be commenced?

When should feeding be commenced?

18. The ESPGHAN expert group recommends to start jejunal feeding within 24 hours after placement of the jejunal feeding tube irrespective of patient age or condition except in complicated surgical situations such as adhesions.

LoE: moderate

SoR: strong

Vote: 9,8,9,9,9,9,9,9,9,5,9,9,9 (93% agreement)

Practical note

Refeeding syndrome should be considered whenever nutritional support is instituted in malnourished children.

Previously, commencement of feeding was delayed until 12 to 24 hours after transabdominal gastrostomy placement to allow the GI tract to return to normal function and to allow healing of the enteral opening. However, several prospective RCTs (90–93) have clearly demonstrated that feeding can be safely started a few hours after the procedure (59), or at least on the first operative day (94), even in early infancy (12).

Abdominal intervention or severe stress are not a contraindication for early feeding as small intestinal motility and absorptive functions have been demonstrated to remain intact, although gastric and colonic motility may be impaired for up to 2 to 5 days (95). Retrospective and prospective observational studies have shown that early transpyloric EN starting within the first 24 hours was well tolerated even in critically ill children without an increased rate of complications compared to late (after 24 hours, range 1–43 days) transpyloric EN (96). Seventy-four percent of the patients achieved their estimated caloric requirements within 24 hours and the remaining patients within 48 hours after transpyloric tube placement (95). Moreover, the incidence of abdominal distension was lower in the children receiving early transpyloric feeding (3.5%) than in those receiving nutrition at a later time (7.8%; $P < 0.05$) (96).

Refeeding syndrome should be considered whenever nutritional support is instituted in malnourished children. It is characterized by electrolyte depletion, fluid shifts, and glucose derangements upon reinstitution of nutrition in malnourished patients.

Q8: Which feeds are suitable for jejunal feeding and what are the nutritional considerations?

19. The ESPGHAN expert group recommends starting feeding with standard polymeric formula, and if this is not tolerated switching to a hydrolysed formula.

LoE: moderate

SoR: strong

Vote: 9,9,9,9,9,9,9,5,9,9,9,6,9,9,9 (87% agreement)

Practical note

Elemental formula and other hyperosmolar feeds should be used with caution.

Thickened and fibre containing feeds should be used with caution due to risk of tube blockage.

20. The ESPGHAN expert group recommends to start with a hydrolysed formula containing medium-chain triglyceride where JTF is used in pancreatic insufficiency or malabsorption.

LoE: low

SoR: strong

Vote: 9,8,9,9,9,9,8,8,8,7,8,8,9,9,9 (100% agreement)

21. The ESPGHAN expert group recommends to monitor serum levels of copper, zinc, selenium, and iron for nutritional deficiencies in all patients who receive long-term JTF.

LoE: low

SoR: strong

Vote: 9,8,7,9,9,9,8,8,8,7,9,9,9,9,9 (100% agreement)

Practical note

Serum levels for copper, zinc, selenium, and iron should be checked on a 6 monthly to 1 yearly basis.

22. The ESPGHAN expert group recommends not to dilute the formulas so as to minimize the risk of microbial contamination of the formula, secondary diarrhoea, and malnutrition due to its low caloric content.

LoE: moderate

SoR: strong

Vote: 9,8,9,9,9,8,9,9,8,9,8,9,9,6,8 (93% agreement)

In children, postpyloric feeds have traditionally been hydrolysed and less viscous because of the narrow lumen of the transpyloric tubes, although polymeric feeds have also been tolerated (97). Evidence in the literature for a particular feed for JTF is however lacking. Physiologically, intraluminal pressure and motility can increase in postpyloric feeding in response to volume and osmolality of the feed. This in turn can cause side effects such as abdominal distension, vomiting, diarrhoea, and dumping syndrome. In postpyloric delivery of feeds pancreatic secretion may vary according to the site and type of feeding. Placement of the tube >40 cm below the ligament of Treitz inhibits pancreatic secretion and this would therefore favour use of an elemental feed. However, O'Keefe et al (98) looked at the effect of polymeric versus an elemental feed on pancreatic secretion. The polymeric feed allowed an adequate pancreatic secretory response, whereas pancreatic secretion was reduced by 50% with the elemental diet. They concluded that intraduodenal infusion allows complete assimilation of a

polymeric enteral feed due to adequate pancreatic secretory response (98).

The underlying disease may also affect the choice of formula, that is, for those with pancreatic insufficiency or malabsorption a semidigested formula may be the feed of choice (99). The higher osmolality of elemental feeds may, however, cause nontolerance. The recommended osmolality for infants and children <4 years of age is <400 mOsm/kg and for older children it is <600 mOsm/kg (100).

It is not recommended to dilute the formulas because it may increase the risk for microbial contamination of the formula (101), secondary diarrhoea, and malnourishment due to its low caloric content (102).

JTF causes an iatrogenic bypass of the upper GI tract, which may lead to nutritional deficiencies. Copper is primarily absorbed in the stomach and therefore in those being jejunally fed there is an increased risk for copper deficiency. Jacobson et al (103) described 3 paediatric patients on exclusive jejunal feeds who developed cytopenia secondary to copper deficiency.

Children on exclusive jejunal feeds may be at risk for iron deficiency due to feeds bypassing the duodenum, which is the primary site for iron absorption. A small case series of 6 children fed via the jejunum showed significant reductions in serum iron (18.5 vs 9.8 g/L, $P = 0.01$) and transferrin levels (23.1% vs 13.7%, $P = 0.02$) after a mean period of 11 months. There was no change in ferritin, haemoglobin, and mean corpuscular volume showing the proximal jejunum may have the capacity to adapt to iron deficiency (104). A retrospective study by Skelton et al (105) showed a 30% reduction in zinc, a 68% reduction in selenium, and a 25% reduction in iron.

There is an increasing popularity amongst families to use blenderized diets in those children on long-term enteral feeds. Blenderized diets need to be given as bolus gravity feed, thereby excluding their use in continuous JTF.

Q9: What feeding regimen should be used for long-term jejunal feeding?

Mode and rates of delivery

23. The ESPGHAN expert group recommends to administer jejunal feeding continuously via a volumetric enteral pump at a rate tailored to the patient's tolerance.

LoE: low

SoR: strong

Vote: 9,9,9,9,7,9,9,8,9,9,8,8,9,9,8 (100% agreement)

There is no evidence indicating the exact rate of the EN delivery. Jejunal feeding should be provided continuously via volumetric enteral pump because bolus feeding or high infusion rate can cause diarrhoea, abdominal cramping, and dumping syndrome-like symptoms (99). Proposed increments are 1 to 5 mL/h every 24 hours for infants or 5 to 20 mL/h in older children every 4 hours until the target rate is reached (9,107).

Suggested volume rates are presented in Table 4. Lower perfusion rates such as $0.5 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{h}^{-1}$ are proposed if there is a risk of gut ischaemia. Once the target rate has been achieved, the concentration of the formula can be increased to deliver the estimated nutrient needs.

Over time, continuous feeding can be cycled with mainly overnight continuous feeding at the highest tolerated rate.

Intermittent continuous feeding is more physiological, allowing greater patient mobility and stimulating oral feeding as it evokes periods of hunger and satiety. Intermittent continuous feeding provides cyclical secretion of GI hormones with a trophic effect on intestinal mucosa (108). Therefore, intermittent continuous feeding patterns would be recommended to use over continuous feeding whenever possible (6).

The rest of the caloric intake can be provided during the day either orally if the child tolerates oral or gastric intake, or via continuous JTF over several hours at the highest tolerated rate. The quantity of feeds per day should be determined by the child's energy requirements and the duration of fasting, which is maximally tolerated. However, JTs should be accessed several times per day even if not in use to maintain tube patency.

When full or partial postpyloric enteral feed cannot be achieved (eg, by clinical instability, airway management, radiological and surgical procedures, and accidental feeding tube removal) (109) *trophic EN* is recommended as continuous infusion of small amounts of enteral feed. Different rates are proposed ranging from 0.5 to $25 \text{ mL} \cdot \text{kg}^{-1} \cdot \text{day}^{-1}$ or 20 mL/h (102,107). Trophic feeding maintains the intestinal barrier and the mucosal integrity and stimulates intestinal secretion of brush border enzymes, endogenous peptides, secretory immunoglobulin A, and bile salts (56,110). These local intestinal effects reduce systemic inflammation by helping to prevent translocation of bacteria or bacterial products across the intestinal epithelial barrier (111).

Q10: What else can the JT be used for?

Administration of medication

24. The ESPGHAN expert group recommends not to use the JTF for the administration of medication unless absolutely essential and/or delivery into the stomach is not possible.

LoE: low

SoR: strong

Vote: 9,9,9,9,7,9,9,7,8,9,8,7,9,9,9 (100% agreement)

Practical note

Delivery of medication via the jejunal route may have unpredictable bioavailability of the active component, the absorption site of the drug, potential local adverse effects, and potential reaction with the feeding tube (possibility of tube clogging).

In the case of motility disorders or obstruction, gastric aspiration should be avoided or performed after a sufficient time interval if medication is administered via the gastric tube.

The jejunal feeding tube can be used for the administration of medications. However, information is lacking regarding the site of intestinal absorption of most medications (112) and drug information sheets usually do not provide information about safety for jejunal intake. Furthermore, in patients with a GJT frequent changes of administration route (gastric vs jejunal) make it difficult to achieve stable therapeutic drug levels (112). Therefore, clinicians have to closely monitor medical therapy in these children. In patients with motility disorders and obstructions, the gastric port of the GJT is sometimes used to aspirate gastric contents. In this case, administration of medication through the gastric or JT should be avoided or at least be performed after a

TABLE 4. Infusion rates based on Pedrón Giner et al (102) and NASPGHAN (106)

Age	Initial phase rate	Advance rate	Suggested tolerated rate
Preterm	0.5–2 mL · kg ⁻¹ · h ⁻¹	0.2–1 mL/kg every 8 h	4–8 mL · kg ⁻¹ · h ⁻¹
Infant	1–2 mL · kg ⁻¹ · h ⁻¹	1–2 mL/kg every 2–8 h	5–6 mL · kg ⁻¹ · h ⁻¹
1–6 y	1 mL · kg ⁻¹ · h ⁻¹	1 mL/kg every 2–8 h	1–5 mL · kg ⁻¹ · h ⁻¹
≥7 y	25 mL/h	25 mL every 2–8 h	100–150 mL/h

sufficient interval, as the medication may be sucked before intestinal absorption (112).

As the JT is often both used for EN and medication, interactions between the food and medication are possible, but there is limited published data in children (112). Additional adverse drug reactions are possible. Many liquid formulations of medications have a high osmolality, which can result in cramping, abdominal distension, vomiting, and diarrhoea (113). These symptoms are usually attributable to inactive ingredients and excipients in the drug formulation, such as, for example, polyethylene glycol (114) or sorbitol (115).

Alterations in drug absorption can lead to increased toxicity or treatment failure (112). Increased toxicity may occur due to a lack of degradation by stomach acid or decreased first-pass hepatic metabolism, leading to increased drug absorption and/or greater systemic exposure (115,116). On the contrary, treatment failure may be caused by decreased absorption time leading to impaired degradation of medication (112,116). Furthermore, bypassing the stomach may reduce the absorption and degradation of pH-dependent drugs (116).

In a large literature review 70 medications had information available regarding GI site of absorption (112) (Table 5).

For the majority of medications, there are no specific data on bioavailability or solubility after the drug (tablet or capsule) is crushed. Many compounds are water-insoluble, and sustained- or extended-release product formulations should not be crushed due to potential toxicity from the rapid release of large doses of the active component (116).

Ideally, to prevent jejunal feeding tube blockage, medications should be completely dissolved in water or applied as liquid formulations (59,117). After administration flushing the tube with water helps to deliver the drug to the intestinal mucosa (118).

Gastric decompression and aspiration

25. The ESPGHAN expert group recommends to perform gastric decompression and aspiration in children being fed by jejunal feeding tube who have a high risk of gastroesophageal reflux and pulmonary aspiration due to accumulation of gastric residue and abdominal distension.

LoE: low

SoR: strong

Vote: 9,8,9,9,9,9,9,8,8,9,8,6,9,9,9 (93% agreement).

Gastrojejunal feeding is a well-established feeding method to provide both postpyloric feeding and gastric decompression in patients with a high risk for GOR and pulmonary aspiration due to accumulation of gastric residue and abdominal distension (75,89,119). In children with an NJT or a surgical jejunostomy

the presence of a gastric tube or a PEG may reduce the risk for GOR and pulmonary aspiration by facilitating gastric decompression.

“Downstream” JTF increases gastric acid secretions (89). Furthermore, there is retrograde movement of enteral feed and bile into the stomach due to abnormal GI motility (89,120). Increased gastric residue/ aspirates are considered as >50% of the volume administered in the previous 4 hours (121). Increase in gastric residue leads to a higher risk of aspiration and also favours SIBO (121).

Aspiration of gastric residue can also help in deciding when to start and how to advance oral feeding when no biliary drainage exists in the nasogastric aspirate (37).

SPECIAL USE IN THE CASE OF SUSPECTED SMALL INTESTINAL BACTERIAL OVERGROWTH

The jejunal feeding tube offers a unique opportunity to aspirate intestinal fluid and evaluate for SIBO in the case of clinical suspicion (eg, bloating, diarrhoea, growth failure) and/or diagnostic indications (vitamin B12 deficiency, urinary organic acids profile). Ideally, aspiration of jejunal secretions for culture should be performed via a new JT to avoid culturing bacteria that have been colonizing the tube. The presence of a (polyethylene) JT itself has

TABLE 5. Information on the absorption site of a list of medications according to McIntyre (112)

Drugs requiring acid for absorption	Aspirin
	Ferrous sulphate
Drugs that bind extensively to the tube	Cyclosporine
	Isotretinoin
Drugs with higher absorption rate when administered in the small bowel	Azathioprine
	Ciprofloxacin
	Fluconazole
	Pravastatin
	Zinc
Drugs with decreased absorption when administered in the small bowel	Allopurinol
	Baclofen
	Calcium
	Ferrous sulphate
	Gabapentin
	Lopinavir
	Ritonavir
	Sirolimus
Drugs not absorbed when administrated in the jejunum	Digoxin
	Erythromycin
	Folic acid
	Griseofulvin
	Metformin
	Mycophenolate
	Phenytoin
	Pravastatin

shown to alter the intestinal flora of the small intestine in very low birth weight preterm infants (122). There was an increased risk to develop NEC if the jejunum was heavily colonized with Gram-negative bacilli (122). Furthermore, increase in gastric residues favours SIBO (121).

Q11: What is needed for on-going care of postpyloric feeding?

26. The ESPGHAN expert group recommends to tailor the care and management of jejunal feeding devices according to the type of device used and route of insertion.

LoE: low

SoR: strong

Vote: 9,8,3,9,9,9,9,8,9,8,7,9,9,7 (93% agreement)

NASOJEJUNAL TUBES

NJTs should be measured before use to rule out displacement. The nasal passage should be clean and dry with good skin integrity. It should then be secured with appropriate tape to avoid re-passing.

To reduce the risk of pulmonary aspiration, the patient should ideally be nursed at a 30° angle or higher if possible.

NASAL BRIDLE RETAINING SYSTEM

Patients requiring NJT placement may benefit from a nasal bridle tube-retaining system. Nasoenteric feeding tubes can become dislodged due to patient noncompliance, transfers, or positional changes (123). Nasal bridles can provide a better, more reliable system to secure nasoenteric tubes.

Placement of a nasal bridle retaining system should only be carried out by health care professionals with specific training.

Contraindications for the insertion of a nasal bridle system include a grossly deviated nasal septum and persistent vomiting (124). Consideration for insertion includes nasal polyps, nasal deformity, a history of epistaxis, and ethnic/cultural issues.

ROUTINE CARE OF SURGICALLY AND ENDOSCOPICALLY PLACED JEJUNAL TUBE

Appropriate labelling should be used for PEG-J tubes distinguishing the gastric and jejunal lumina. Depending on the manufacturer some devices have a balloon-retaining bumper, which requires weekly water changes to make sure the balloon is always filled. GJTs are not to be rotated to avoid migration back into the stomach. Feeding extension sets are to be changed as per manufacturer's instructions. Minimal handling and an aseptic nontouch technique should always be applied to connect the administration set to the enteral feeding tube and feed receptacle (125). The exit site is cleaned and dried at least once a day. PEJ/PEG-J tubes do not require routine aspiration but if the patient is showing signs of respiratory distress or vomiting then the pH of aspirate should be checked. A pH <5.5 may indicate that the tube has migrated to the stomach and the feed should be stopped and the tube checked with an x-ray (126). If an aspirate were obtainable from the jejunum then a pH of 6 to 8 would be expected (127).

Tube flushing

27. The ESPGHAN expert group recommends to flush the jejunal feeding tube with small amounts of warm water before and after administration of EN and medication or when changing the bag or bottle in the case of continuous JTF.

LoE: very low

SoR: strong

Vote: 9,9,7,9,9,8,9,6,7,7,9,9,9,9 (93% agreement)

To maintain patency, the tube should be flushed whenever the feed is interrupted, before and after all feeds and medication administration (102) with 10 to 20 mL of sterile water 4 to 8 hourly (unless the child is fluid restricted). The water must be sterile.

All medications should be administered in liquid form; some liquid medications are known to be associated with tube blockages and so can be diluted before administering via the tube.

If giving 2 or more medications at the same time flushing is recommended in between to prevent precipitation/ clogging (settling of the medication) in the tubing.

TUBE BLOCKAGE

Mechanical complications are frequently reported (10–12); and often related to inadequate tube care by caregivers and nursing staff (59). Such events may be prevented by correct education on tube management with the goal to avoid frequent tube substitution. Several agents have been proposed for prevention and treatment of tube clogging, including pancreatic enzymes and carbonated beverages, but in vivo trials are still needed to establish their efficacy (128). Flushing may be more effective with warm water and small-volume syringes (1, 2, or 5 mL) to create higher pressure.

SITE MANAGEMENT

Feed Handling and Preparation

In recent years, powdered infant formula contaminated with harmful bacteria has been associated with serious illness and death due to infection with bacteria such as *Cronobacter sakazakii* (129). Following this, recommendations on preparation of powdered infant formula have changed both for parents at home and in health care settings (Table 6). The World Health Organization guideline on "Safe Preparation, Storage and Handling of Powdered Infant Formula" (130) states that: "Powdered and decanted liquid feeds should only be used when there is no suitable alternative sterile feed available."

The handling of the enteral feed should be done in a clean environment using aseptic techniques by trained staff and if required the feed should be reconstituted with sterile or purified water heated to 70°C to 80°C (102). A prolonged hanging time increases the risk for retrograde contamination and, therefore, the hanging time should not exceed beyond 24 hours (129,131). Feed continuously administered should not be warmed.

Storage of feeds

28. The ESPGHAN expert group recommends to use a closed system for the preparation of the feed to avoid infection and error (eg, correct feed, use before expiry date).

LoE: very low

SoR: strong

Vote: 9,7,8,9,9,9,9,8,9,9,8,9,9,9 (100% agreement)

TABLE 6. Site management of jejunal tubes

Stoma site appearance	Treatment/management
Healthy stoma	Stoma site should be clean and dry. It should be daily cleaned when bathing the child and dried thoroughly
Redness to site	Assess patient, consider cellulitis/collection. Consider tube size. If there is moisture consider foam dressings and barrier cream. Swab stoma and send to microbiology. Consider topical creams for inflammation.
Discharge to site	Low to moderate exudate: thin foam dressing.
Assess and document redness	High exudate: apply absorbent foam antimicrobial dressings. Apply barrier cream to protect skin. Assess tube and fit. Send swab for microbiology. Consider oral antibiotic.
Leakage from stoma site	Try venting the stomach to relieve pressure. Thoroughly clean and dry stoma site. Cover with nonadhesive foam dressings. If leakage persists and irritates the skin apply barrier creams to protect the skin and continue to cover with foam dressing.
Granulation tissue (pink, moist tissue around stoma, easily bleeding) Causes:	Prevent by securing tube with tape. Check if fixation device is in correct position and fits well. Avoid friction to the site. Foam dressings on small granulation.
- friction (seat belts/ clothing)	Topical ointments for up to 7 days for moderate granulation.
- tube pulled too tightly or excessive movement of device	Silver Dressings or silver nitrate for persistent granulation tissue (by experienced practitioner).
- bacterial colonisation of site causing inflammation	In rare cases surgical therapy is sought for huge overgranulation that has failed medical management.

The World Health Organization recommends storage of feeds in a clean, dark place in its original box, between 15°C and 25°C avoiding extreme temperatures, to avoid handling whenever possible (102,130). Prepared feeds should not be frozen. The feed must always be connected to the administration set according to the manufacturer's instructions, and always with an aseptic nontouch technique (132,133). In bedded services the feeding system must always be labelled with the patient's name and the date and time the feed was set up (134). All opened containers of ingredients should be covered, labelled with an expiry date, and stored in a clean secure location. Dry ingredients once opened should be used within 4 weeks of opening or as determined by the manufacturers' instructions if sooner. All opened or unused made-up liquid feeds must be discarded in accordance to the manufacturer's instructions.

Home enteral nutrition (HEN) is now widely supported and recognized as beneficial for the child's well-being and maintaining

the family unit (135). Communication between hospital and community MDT (ideally consisting of a paediatric gastroenterologist, a dietician, a psychologist, an occupational therapist, and a speech therapist) involved in the child's care is therefore paramount between primary and secondary care settings in providing safe and effective care (136).

In hospital settings, education and training is provided to parents/caregivers on preparation of feeds, management of feeding device and skin sites, pump training, and administration of nutrition and medications. Various members of the MDT are involved throughout all stages of training (nurses, dietician and medical doctors) (137). Communication and support to families is paramount as it offers good understanding, provides safety and equips families with a competent feeling to take over care once discharged home (136).

The child's primary care giver follows up the child after JT insertion with the help of community nurses, dieticians and at times speech and language therapists. This provides a more holistic care approach making sure that the child is thriving well as that the tube stoma tract is being well maintained. This shows that even in the community setting an MDT approach is also essential.

A number of hospital centres/ teams have developed tube feeding clinics to offer a holistic approach of care for these highly complex chronically ill children (138). These teams generally include a physician (usually a paediatric gastroenterologist), a dietician, a nutrition nurse, a speech and language therapist, and parents/caregivers (132,133).

Feeding clinics focus mainly on addressing, restoring and maintaining an adequate nutritional status of children to avoid nutritional depletion and to allow children to reach their potential growth and development. The timing of each child's feeding clinic

Q12: Who should be involved in the follow-up care?

29. The ESPGHAN expert group recommends to use an MDT approach with well trained professionals for the follow-up and management of children requiring jejunal feeding.

LoE: low

SoR: strong

Vote: 9,9,9,9,9,9,8,9,9,9,9,9,9 (100% agreement)

review therefore varies depending on its nutritional status and general well-being.

Q15: How should you wean off jejunal tube feeding?

30. The ESPGHAN expert group recommends, in the absence of a standard approach such as a clinical guideline, to wean off JTF using an MDT setting providing an on-going monitoring and support.

LoE: low

SoR: strong

Vote: 9,9,9,9,9,9,8,9,9,9,9,9,9 (100% agreement)

Practical note

If possible children on JTF should be re-trialled on oral or gastric feeds at intervals.

There is little guidance in the literature as to how to wean successfully off a JT (139,140). As with all tube feeding when a tube is placed there should always be a discussion around transitioning back from tube to oral feeding where possible. The decision to begin weaning the child from JTF will depend on nutritional status, medical stability, and oral aversion. Transition may take days to months and depending on the reason for the JTF a child may remain dependent on the JT for a long period of time. Feeding aversion may be a major issue and where possible some continuation of oral (if safe) feeding should be considered to limit this aversion.

Transition of JTF back to gastric/oral feeding may be achieved by many methods and will depend on a centre's practice (141). The child may be admitted to hospital or a specialist centre for tube weaning or the child may remain at home with small changes made over a period of time. Interventions consist of psychoeducation, supportive psychotherapy for parents including parent-child relationship work, behavioural interventions with mealtime structuring, nutritional and medical interventions, hunger provocation, and treatment of oral sensory-motor difficulties (142). Where possible tube weaning should be done within the setting of an MDT comprising of a dietician, specialist speech and language therapist, psychologist, occupational therapist, and a paediatrician.

Q16: What are the ethical considerations?

31. The ESPGHAN expert group recommends to involve parents and/ or caregivers in each decision-making process and to ensure that informed consent is obtained.

LoE: moderate

SoR: strong

Vote: 9,8,8,9,9,9,8,9,9,9,9,9,8 (100% agreement)

32. The ESPGHAN expert group recommends to involve a professional ethicist to assist in decision-making in cases where the insertion of a jejunostomy poses ethical dilemmas.

LoE: very low

SoR: strong

Vote: 9,8,9,9,9,9,7,8,9,8,9,9,9 (100% agreement)

JTF is a therapeutic intervention aiming at reversing malnutrition and/or maintaining nutritional status in children who cannot tolerate oral or gastrostomy feeds. Therefore, similar to other diagnostic or therapeutic interventions the decision on its initiation must be a result of consensus between the medical professionals and the parents/caregivers. Obtaining informed and educated consent by the parents/caregivers is an important ethical principle of every invasive intervention procedure including JTF.

The benefit of JTF is determined by the potential medical benefits but also by the perceived benefits by the child's parents/caregivers. Sometimes, the decision-making process regarding tube feeding for parents/caregivers is difficult and the process is delayed (143,144). Multiple negative perceptions may coexist including feeling of failure, disruption of maternal nurturing and bonding, loss of normality, and confirmation of the permanence of the disability. It is therefore important for the MDT to recognise these perceptions and to be involved in the discussions with the parents/caregivers explaining benefits, risks, alternatives, and the consequences of not receiving the proposed treatment. The parents/caregivers should be given enough time to make their decision freely. The ethical principle of informed consent is based on the understanding of the above perceptions by the health care professionals (145) who also need to develop effective, family-centred, patient-appropriate adherence strategies.

Furthermore, to promote the best interest of the patient it is important to make every effort to guarantee maximum effectiveness of the intervention with minimum complications at reasonable costs. To achieve the above goals, the jejunal feeding tubes should be placed by experienced specialists and the jejunal feeding should be supervised by specifically trained professionals. Care coordination by an MDT including the families/caregivers, improves outcomes in patients receiving long-term enteral feeding, whereas specialized home enteral tube feeding programs significantly reduce morbidity and costs (146,147). The ESPGHAN Committee on Nutrition recommends the implementation in hospitals of multidisciplinary nutritional care teams with expertise in all aspects of clinical nutrition care, funded by the health care system (138).

In conclusion, the decision to establish JTF must be based on the best clinical evidence and take into consideration the clinician's experience and the parents'/caregivers' perceptions, concerns, and expectations. Acknowledgement of benefits, risks, costs, and effects in the decision-making process provide the best approach for both health professionals and parents/caregivers ultimately promoting the patient's optimal growth, health, and quality of life.

Q17: Who is involved in the management at home/in the community?

33. The ESPGHAN expert group recommends that in all patients on HEN there is close cooperation between the home (parents/caregivers and community nursing team) and hospital MDTs.

LoE: low

SoR: strong

Vote: 9,9,8,9,9,9,8,9,9,9,9,9,8 (100% agreement)

HEN provides nutritional support to children with chronic diseases allowing them to be discharged earlier from hospitals (102). Only a minority of patients receive their feeding via jejunal approach (148), and general aspects of HEN may be applied to this subgroup. Data from a Spanish national paediatric registry on 952

patients on HEN show that the majority of patients are fed enterally due to neurological disease. The number of HEN patients has increased substantially over 1 decade. However, only 2.2% of patients in the registry are fed jejunally (102). There are no RCTs available on home/community involvement in the care of patients with jejunal feeding.

When planning for discharge several important factors must be considered, namely stability of the patient's condition, adequate psychomotor skills, and ability to understand and retain information. Adequate education and training, as well as supply of all necessary equipment required for HEN is essential. Referral to respective specialists (dietician, general practitioner or general paediatrician, gastroenterologist, etc) able to prescribe the feeds in the outpatient setting needs to be ensured (6,135,148). In some countries commercial feeding companies can provide training for patients, caregivers, including, for example, "out of hours" advice lines where patients and caregivers can obtain troubleshooting information (148). As community follow-up is often inadequate for patients discharged home on enteral tube feeding (136,137,148) and poor discharge information leads to predominantly negative experience of general paediatricians with enteral feeding (137), optimal communication at discharge between health care professionals in secondary and primary care services and MDT needs to be established (6,132,133,135,148–150). Close cooperation should be established also between the community and the hospital nutrition team, if available. The need for hospital nutrition teams has been stressed by the ESPGHAN Committee on Nutrition (138).

Essential information given to the parents or caregivers and possibly to children at discharge should include reasons for home tube feeding and likely duration; safety aspects of care (tube placement, infection control, hand-washing, feed preparation); information on feeding equipment; social and practical implications; problem-solving advice; the importance of maintaining oral stimulation; telephone contacts for hospital and community staff; and detailed information about how to obtain equipment and supplies (6,135,148). The use of an easy-to-manage, lightweight, and portable enteral feeding pump is recommended for jejunal feeding and detailed instructions on the management of the pump should be given at discharge (6). Information on regular evaluation of the nutritional status and oral motor skills, swallowing, and gastroesophageal function is essential to allow early taper of jejunal EN (6,149).

ESPGHAN disclaimer:

ESPGHAN is not responsible for the practices of physicians and provides guidelines and position papers as indicators of best practice only. Diagnosis and treatment are at the discretion of physicians.

Acknowledgments: The authors thank Giulia Angelino for the preparation of Table 3 and review of the literature to questions 5 and 6.

REFERENCES

- Campwala I, Perrone E, Yanni G, et al. Complications of gastrojejunal feeding tubes in children. *J Surg Res* 2015;199:67–71.
- Egnell C, Eksborg S, Grahngquist L. Jejunostomy enteral feeding in children: outcome and safety. *JPEN J Parenter Enteral Nutr* 2014;38:631–6.
- King M, Barnhart DC, O'Gorman M, et al. Effect of gastrojejunal feedings on visits and costs in children with neurologic impairment. *J Pediatr Gastroenterol Nutr* 2014;58:518–24.
- Rosen R, Vandenplas Y, Singendonk M, et al. Pediatric Gastroesophageal Reflux Clinical Practice Guidelines: Joint Recommendations of the North American Society for Pediatric Gastroenterology, Hepatology, and Nutrition and the European Society for Pediatric Gastroenterology, Hepatology, and Nutrition. *J Pediatr Gastroenterol Nutr* 2018;66:516–54.
- Romano C, Van Wynckel M, Hulst J, et al. European Society for Paediatric Gastroenterology, Hepatology and Nutrition guidelines for the evaluation and treatment of gastrointestinal and nutritional complications in children with neurological impairment. *J Pediatr Gastroenterol Nutr* 2017;65:242–64.
- Braegger C, Decsi T, Dias JA, et al. Practical approach to paediatric enteral nutrition: a comment by the ESPGHAN Committee on Nutrition. *J Pediatr Gastroenterol Nutr* 2010;51:110–22.
- Thapar N, Saliakellis E, Benninga MA, et al. Paediatric intestinal pseudo-obstruction: evidence and consensus-based recommendations from an ESPGHAN-Led Expert Group. *J Pediatr Gastroenterol Nutr* 2018;66:991–1019.
- McClave SA, Taylor BE, Martindale RG, et al. Guidelines for the provision and assessment of nutrition support therapy in the adult critically ill patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J Parenter Enteral Nutr* 2016;40:159–211.
- Mehta NM, Compheer C. Directors ASPENBo A.S.P.E.N.. Clinical guidelines: nutrition support of the critically ill child. *JPEN J Parenter Enteral Nutr* 2009;33:260–76.
- Godbole P, Margabanthu G, Crabbe DC, et al. Limitations and uses of gastrojejunal feeding tubes. *Arch Dis Child* 2002;86:134–7.
- Fortunato JE, Darbari A, Mitchell SE, et al. The limitations of gastrojejunal (G-J) feeding tubes in children: a 9-year pediatric hospital database analysis. *Am J Gastroenterol* 2005;100:186–9.
- Michaud L, Coopman S, Guimber D, et al. Percutaneous gastrojejunostomy in children: efficacy and safety. *Arch Dis Child* 2012;97:733–4.
- Guyatt GH, Oxman AD, Vist GE, et al. GRADE: an emerging consensus on rating quality of evidence and strength of recommendations. *BMJ* 2008;336:924–6.
- Guyatt GH, Oxman AD, Kunz R, et al. GRADE guidelines: 2. Framing the question and deciding on important outcomes. *J Clin Epidemiol* 2011;64:395–400.
- Balshem H, Helfand M, Schunemann HJ, et al. GRADE guidelines: 3. Rating the quality of evidence. *J Clin Epidemiol* 2011;64:401–6.
- Guyatt GH, Oxman AD, Vist G, et al. GRADE guidelines: 4. Rating the quality of evidence—study limitations (risk of bias). *J Clin Epidemiol* 2011;64:407–15.
- Hsu J, Brozek JL, Terracciano L, et al. Application of GRADE: making evidence-based recommendations about diagnostic tests in clinical practice guidelines. *Implement Sci* 2011;6:62.
- Guyatt G, Oxman AD, Akl EA, et al. GRADE guidelines: 1. Introduction—GRADE evidence profiles and summary of findings tables. *J Clin Epidemiol* 2011;64:383–94.
- Alkhawaja S, Martin C, Butler RJ, et al. Post-pyloric versus gastric tube feeding for preventing pneumonia and improving nutritional outcomes in critically ill adults. *Cochrane Database Syst Rev* 2015;Cd008875.
- Li Z, Qi J, Zhao X, et al. Risk-benefit profile of gastric vs transpyloric feeding in mechanically ventilated patients: a meta-analysis. *Nutr Clin Pract* 2016;31:91–8.
- Martinez EE, Bechard LJ, Mehta NM. Nutrition algorithms and bedside nutrient delivery practices in pediatric intensive care units: an international multicenter cohort study. *Nutr Clin Pract* 2014;29:360–7.
- Zhang Z, Xu X, Ding J, et al. Comparison of postpyloric tube feeding and gastric tube feeding in intensive care unit patients: a meta-analysis. *Nutr Clin Pract* 2013;28:371–80.
- Meert KL, Daphtary KM, Metheny NA. Gastric vs small-bowel feeding in critically ill children receiving mechanical ventilation: a randomized controlled trial. *Chest* 2004;126:872–8.
- Watson J, McGuire W. Transpyloric versus gastric tube feeding for preterm infants. *Cochrane Database Syst Rev* 2013;CD003487.
- Di Lorenzo C, Flores AF, Buie T, et al. Intestinal motility and jejunal feeding in children with chronic intestinal pseudo-obstruction. *Gastroenterology* 1995;108:1379–85.
- Saliakellis E, Fotoulaki M. Gastroparesis in children. *Ann Gastroenterol* 2013;26:204–11.
- Tillman EM, Smetana KS, Bantu L, et al. Pharmacologic treatment for pediatric gastroparesis: a review of the literature. *J Pediatr Pharmacol Ther* 2016;21:120–32.

28. Camilleri M, Parkman HP, Shafi MA, et al. Clinical guideline: management of gastroparesis. *Am J Gastroenterol* 2013;108:18–37quiz 38.
29. Goulet O, Olieman J, Ksiazyk J, et al. Neonatal short bowel syndrome as a model of intestinal failure: physiological background for enteral feeding. *Clin Nutr* 2013;32:162–71.
30. Tappenden KA. Intestinal adaptation following resection. *JPEN J Parenter Enteral Nutr* 2014;38(1 suppl):23s–31s.
31. Livingston MH, Shawyer AC, Rosenbaum PL, et al. Fundoplication and gastrostomy versus percutaneous gastrojejunostomy for gastroesophageal reflux in children with neurologic impairment: a systematic review and meta-analysis. *J Pediatr Surg* 2015;50:707–14.
32. Srivastava R, Downey EC, O’Gorman M, et al. Impact of fundoplication versus gastrojejunal feeding tubes on mortality and in preventing aspiration pneumonia in young children with neurologic impairment who have gastroesophageal reflux disease. *Pediatrics* 2009;123:338–45.
33. Wales PW, Diamond IR, Dutta S, et al. Fundoplication and gastrostomy versus image-guided gastrojejunal tube for enteral feeding in neurologically impaired children with gastroesophageal reflux. *J Pediatr Surg* 2002;37:407–12.
34. Albanese CT, Towbin RB, Ulman I, et al. Percutaneous gastrojejunostomy versus Nissen fundoplication for enteral feeding of the neurologically impaired child with gastroesophageal reflux. *J Pediatr* 1993;123:371–5.
35. Rosen R, Hart K, Warlaumont M. Incidence of gastroesophageal reflux during transpyloric feeds. *J Pediatr Gastroenterol Nutr* 2011;52:532–5.
36. Stone B, Hester G, Jackson D, et al. Effectiveness of fundoplication or gastrojejunal feeding in children with neurologic impairment. *Hosp Pediatr* 2017;7:140–8.
37. Jiang W, Lv X, Xu X, et al. Early enteral nutrition for upper digestive tract malformation in neonate. *Asia Pac J Clin Nutr* 2015;24:38–43.
38. Jiang W, Zhang J, Geng Q, et al. Early enteral nutrition in neonates with partial gastrectomy: a multi-center study. *Asia Pac J Clin Nutr* 2016;25:46–52.
39. Abu-El-Haija M, Kumar S, Quiros JA, et al. The management of acute pancreatitis in the pediatric population: a clinical report from the NASPGHAN Pancreas Committee. *J Pediatr Gastroenterol Nutr* 2018;66:159–76.
40. Meyer R, Foong RX, Thapar N, et al. Systematic review of the impact of feed protein type and degree of hydrolysis on gastric emptying in children. *BMC Gastroenterol* 2015;15:137.
41. Dutta S, Singh B, Chessell L, et al. Guidelines for feeding very low birth weight infants. *Nutrients* 2015;7:423–42.
42. Ng E, Shah VS. Erythromycin for the prevention and treatment of feeding intolerance in preterm infants. *Cochrane Database Syst Rev* 2008;Cd001815.
43. Ericson JE, Arnold C, Cheeseman J, et al. Use and safety of erythromycin and metoclopramide in hospitalized infants. *J Pediatr Gastroenterol Nutr* 2015;61:334–9.
44. Ng PC. Erythromycin as a prokinetic agent in newborns—useful or doubtful. Commentary on Y. Mansi et al.: Randomized controlled trial of a high dose of oral erythromycin for the treatment of feeding intolerance in preterm infants (Neonatology 2011;100:290–294). *Neonatology* 2011;100:297–8.
45. Hyman PE, Abrams CE, Dubois A. Gastric emptying in infants: response to metoclopramide depends on the underlying condition. *J Pediatr Gastroenterol Nutr* 1988;7:181–4.
46. Gariepy CE, Mousa H. Clinical management of motility disorders in children. *Semin Pediatr Surg* 2009;18:224–38.
47. Schey R, Saadi M, Midani D, et al. Domperidone to treat symptoms of gastroparesis: benefits and side effects from a large single-center cohort. *Dig Dis Sci* 2016;61:3545–51.
48. Tang DM, Friedenberg FK. Gastroparesis: approach, diagnostic evaluation, and management. *Dis Mon* 2011;57:74–101.
49. Van Haren FM, Ram S, Hill J. Endoscopic findings during placement of postpyloric feeding tubes in intensive care patients: a retrospective observational study. *Crit Care Resusc* 2012;14:101–4.
50. O’Keefe SJ, Foody W, Gill S. Transnasal endoscopic placement of feeding tubes in the intensive care unit. *JPEN J Parenter Enteral Nutr* 2003;27:349–54.
51. Schwab D, Muhldorfer S, Nusko G, et al. Endoscopic placement of nasojejunal tubes: a randomized, controlled, prospective trial comparing suitability and technical success for two different tubes. *Gastrointest Endosc* 2002;56:858–63.
52. McGuire W, McEwan P. Systematic review of transpyloric versus gastric tube feeding for preterm infants. *Arch Dis Child Fetal Neonatal Ed* 2004;89:F245–8.
53. McGuire W, McEwan P. Transpyloric versus gastric tube feeding for preterm infants. *Cochrane Database Syst Rev* 2002;3:CD003487.
54. McGuire W, McEwan P. Transpyloric versus gastric tube feeding for preterm infants. *Cochrane Database Syst Rev* 2007;3:CD003487.
55. Wallenstein MB, Stevenson DK. Need for reassessment of early transpyloric feeding in preterm infants. *JAMA Pediatr* 2018;172:1004–5.
56. Ohta K, Omura K, Hirano K, et al. The effects of an additive small amount of a low residual diet against total parenteral nutrition-induced gut mucosal barrier. *Am J Surg* 2003;185:79–85.
57. Tyson JE, Kennedy KA. Trophic feedings for parenterally fed infants. *Cochrane Database Syst Rev* 2005;3:CD000504.
58. Patchell CJ, Anderton A, MacDonald A, et al. Bacterial contamination of enteral feeds. *Arch Dis Child* 1994;70:327–30.
59. Blumenstein I, Shastri YM, Stein J. Gastroenteric tube feeding: techniques, problems and solutions. *World J Gastroenterol* 2014;20:8505–24.
60. Lyons KA, Brilli RJ, Wieman RA, et al. Continuation of transpyloric feeding during weaning of mechanical ventilation and tracheal extubation in children: a randomized controlled trial. *JPEN J Parenter Enteral Nutr* 2002;26:209–13.
61. Chatwin M, Bush A, Macrae DJ, et al. Risk management protocol for gastrostomy and jejunostomy insertion in ventilator dependent infants. *Neuromuscul Disord* 2013;23:289–97.
62. Roeder BE, Said A, Reichelderfer M, et al. Placement of gastrostomy tubes in patients with ventriculoperitoneal shunts does not result in increased incidence of shunt infection or decreased survival. *Dig Dis Sci* 2007;52:518–22.
63. Khlevner J, Antino J, Panesar R, et al. Establishing early enteral nutrition with the use of self-advancing postpyloric feeding tube in critically ill children. *JPEN J Parenter Enteral Nutr* 2012;36:750–2.
64. October TW, Hardart GE. Successful placement of postpyloric enteral tubes using electromagnetic guidance in critically ill children. *Pediatr Crit Care Med* 2009;10:196–200.
65. Kline AM, Sorce L, Sullivan C, et al. Use of a noninvasive electromagnetic device to place transpyloric feeding tubes in critically ill children. *Am J Crit Care* 2011;20:453–9, quiz 60.
66. Da Silva PS, Paulo CS, De Oliveira Iglesias SB, et al. Bedside transpyloric tube placement in the pediatric intensive care unit: a modified insufflation air technique. *Intensive Care Med* 2002;28:943–6.
67. Tiancha H, Jiyong J, Min Y. How to promote bedside placement of the postpyloric feeding tube: a network meta-analysis of randomized controlled trials. *JPEN J Parenter Enteral Nutr* 2015;39:521–30.
68. Clifford P, Ely E, Heimall L. Bedside placement of the postpyloric tube in infants. *Adv Neonatal Care* 2017;17:19–26.
69. Cone LC, Gilligan MF, Kagan RJ, et al. Enhancing patient safety: the effect of process improvement on bedside fluoroscopy time related to nasoduodenal feeding tube placement in pediatric burn patients. *J Burn Care Res* 2009;30:606–11.
70. Dimand RJ, Veereman-Wauters G, Braner DA. Bedside placement of pH-guided transpyloric small bowel feeding tubes in critically ill infants and small children. *JPEN J Parenter Enteral Nutr* 1997;21:112–4.
71. Krafte-Jacobs B, Carver J, Wilkinson JD. Comparison of gastric intramucosal pH and standard perfusional measurements in pediatric septic shock. *Chest* 1995;108:220–5.
72. Karabulut R, Turkylmaz Z, Sonmez K, et al. A very feasible alternative in patients with feeding difficulties from gastrostomy: jejunal tube advanced through the gastrostomy. *Afr J Paediatr Surg* 2015;12:119–21.
73. Sy K, Dipchand A, Atenafu E, et al. Safety and effectiveness of radiologic percutaneous gastrostomy and gastro jejunostomy in children with cardiac disease. *AJR Am J Roentgenol* 2008;191:1169–74.

74. Raval MV, Phillips JD. Optimal enteral feeding in children with gastric dysfunction: surgical jejunostomy vs image-guided gastrojejunal tube placement. *J Pediatr Surg* 2006;41:1679–82.
75. Picoraro JA, Pierog A, Reilly NR, et al. Gastrojejunal tube placement through an established gastrostomy via an endoscopic transgastric approach in a pediatric population. *Gastrointest Endosc* 2015;82:1025–30.
76. Jacob A, Delesalle D, Coopman S, et al. Safety of the one-step percutaneous endoscopic gastrostomy button in children. *J Pediatr* 2015;166:1526–8.
77. Michaud L, Robert-Dehault A, Coopman S, et al. One-step percutaneous gastrojejunostomy in early infancy. *J Pediatr Gastroenterol Nutr* 2012;54:820–1.
78. Virnig DJ, Frech EJ, Delegege MH, et al. Direct percutaneous endoscopic jejunostomy: a case series in pediatric patients. *Gastrointest Endosc* 2008;67:984–7.
79. Belsha D, Thomson M, Dass DR, et al. Assessment of the safety and efficacy of percutaneous laparoscopic endoscopic jejunostomy (PLEJ). *J Pediatr Surg* 2016;51:513–8.
80. Esposito C, Settini A, Centonze A, et al. Laparoscopic-assisted jejunostomy: an effective procedure for the treatment of neurologically impaired children with feeding problems and gastroesophageal reflux. *Surg Endosc* 2005;19:501–4.
81. Onwubiko C, Bairdain S, Murphy AJ, et al. Laparoscopic gastrojejunostomy tube placement in infants with congenital cardiac disease. *J Laparoendosc Adv Surg Tech A* 2015;25:1047–50.
82. Neuman HB, Phillips JD. Laparoscopic Roux-en-Y feeding jejunostomy: a new minimally invasive surgical procedure for permanent feeding access in children with gastric dysfunction. *J Laparoendosc Adv Surg Tech A* 2005;15:71–4.
83. Schlager A, Arps K, Siddharthan R, et al. The “omega” jejunostomy tube: a preferred alternative for postpyloric feeding access. *J Pediatr Surg* 2016;51:260–3.
84. Jester I, Singh M, Parikh DH. A novel technique of permanent self-catheterizable feeding jejunostomy. *J Pediatr Surg* 2012;47:812–4.
85. Goring J, Lawson A, Godse A. Are PEGs a risk factor for the buried bumper syndrome? *J Pediatr Surg* 2016;51:257–9.
86. Heuschkel RB, Gottrand F, Devarajan K, et al. ESPGHAN position paper on management of percutaneous endoscopic gastrostomy in children and adolescents. *J Pediatr Gastroenterol Nutr* 2015;60:131–41.
87. Babbitt C. Transpyloric feeding in the pediatric intensive care unit. *J Pediatr Gastroenterol Nutr* 2007;44:646–9.
88. Lopez-Herce J, Santiago MJ, Sanchez C, et al. Risk factors for gastrointestinal complications in critically ill children with transpyloric enteral nutrition. *Eur J Clin Nutr* 2008;62:395–400.
89. Lopez-Herce J, Mencia S, Sanchez C, et al. Postpyloric enteral nutrition in the critically ill child with shock: a prospective observational study. *Nutr J* 2008;7:6.
90. Brown DN, Miedema BW, King PD, et al. Safety of early feeding after percutaneous endoscopic gastrostomy. *J Clin Gastroenterol* 1995;21:330–1.
91. Choudhry U, Barde CJ, Markert R, et al. Percutaneous endoscopic gastrostomy: a randomized prospective comparison of early and delayed feeding. *Gastrointest Endosc* 1996;44:164–7.
92. McCarter TL, Condon SC, Aguilar RC, et al. Randomized prospective trial of early versus delayed feeding after percutaneous endoscopic gastrostomy placement. *Am J Gastroenterol* 1998;93:419–21.
93. Stein J, Schulte-Bockholt A, Sabin M, et al. A randomized prospective trial of immediate vs. next-day feeding after percutaneous endoscopic gastrostomy in intensive care patients. *Intensive Care Med* 2002;28:1656–60.
94. Castle SL, Speer AL, Torres MB, et al. Combined laparoscopic-endoscopic placement of primary gastrojejunal feeding tubes in children: a preliminary report. *J Laparoendosc Adv Surg Tech A* 2013;23:170–3.
95. Chellis MJ, Sanders SV, Webster H, et al. Early enteral feeding in the pediatric intensive care unit. *J Parenter Enteral Nutr* 1996;20:71–3.
96. Sanchez C, Lopez-Herce J, Carrillo A, et al. Early transpyloric enteral nutrition in critically ill children. *Nutrition* 2007;23:16–22.
97. Niv E, Fireman Z, Vaisman N. Post-pyloric feeding. *World J Gastroenterol* 2009;15:1281–8.
98. O’Keefe SJ, Lee RB, Anderson FP, et al. Physiological effects of enteral and parenteral feeding on pancreaticobiliary secretion in humans. *Am J Physiol Gastrointest Liver Physiol* 2003;284:G27–36.
99. Capriati T, Cardile S, Chiusolo F, et al. Clinical management of post-pyloric enteral feeding in children. *Expert Rev Gastroenterol Hepatol* 2015;9:929–41.
100. Vermilyea S, Goh VL. Enteral feedings in children: sorting out tubes, buttons, and formulas. *Nutr Clin Pract* 2016;31:59–67.
101. Bankhead R, Boullata J, Brantley S, et al. Enteral nutrition practice recommendations. *J Parenter Enteral Nutr* 2009;33:122–67.
102. Pedron Giner C, Martinez-Costa C, Navas-Lopez VM, et al. Consensus on paediatric enteral nutrition access: a document approved by SENPE/SEGHP/ANECIPN/SECP. *Nutr Hosp* 2011;26:1–15.
103. Jacobson AE, Kahwash SB, Chawla A. Refractory cytopenias secondary to copper deficiency in children receiving exclusive jejunal nutrition. *Pediatr Blood Cancer* 2017;64:1–4.
104. Tan LZ, Adams SE, Kennedy A, et al. Are children on jejunal feeds at risk of iron deficiency? *World J Gastroenterol* 2015;21:5751–4.
105. Skelton JA, Havens PL, Werlin SL. Nutrient deficiencies in tube-fed children. *Clin Pediatr (Phila)* 2006;45:37–41.
106. NASPGHAN. Pediatric Enteral Nutrition: A Comprehensive Review. <https://www.naspghan.org/files/documents/pdfs/medical-resources/nutrition/Enteral%20Nutrition%20Slide%20Deck%20PDF%20COMPLETE-%20v12Oct2012.pdf>. Accessed January 2, 2018.
107. Mehta NM. Approach to enteral feeding in the PICU. *Nutr Clin Pract* 2009;24:377–87.
108. Jawaheer G, Shaw NJ, Pierro A. Continuous enteral feeding impairs gallbladder emptying in infants. *J Pediatr* 2001;138:822–5.
109. De Oliveira Iglesias SB, Leite HP, Santana e Meneses JF, et al. Enteral nutrition in critically ill children: are prescription and delivery according to their energy requirements? *Nutr Clin Pract* 2007;22:233–9.
110. Bragg LE, Thompson JS, Rikkers LF. Influence of nutrient delivery on gut structure and function. *Nutrition* 1991;7:237–43.
111. Berg RD. Bacterial translocation from the gastrointestinal tract. *J Med* 1992;23:217–44.
112. McIntyre CM, Monk HM. Medication absorption considerations in patients with postpyloric enteral feeding tubes. *Am J Health Syst Pharm* 2014;71:549–56.
113. Estoup M. Approaches and limitations of medication delivery in patients with enteral feeding tubes. *Crit Care Nurse* 1994;14:68–72.
114. Shepherd MF, Felt-Gunderson PA. Diarrhea associated with lorazepam solution in a tube-fed patient. *Nutr Clin Pract* 1996;11:117–20.
115. Beckwith MC, Feddema SS, Barton RG, et al. A guide to drug therapy in patients with enteral feeding tubes: dosage form selection and administration methods. *Hosp Pharm* 2004;39:225–37.
116. Prohaska ES, King AR. Administration of antiretroviral medication via enteral tubes. *Am J Health Syst Pharm* 2012;69:2140–6.
117. Behnken I, Gaschott T, Stein J. [Enteral nutrition: drug administration via feeding tube]. *Z Gastroenterol* 2005;43:1231–41.
118. Sriram K, Jayanthi V, Lakshmi RG, et al. Prophylactic locking of enteral feeding tubes with pancreatic enzymes. *J Parenter Enteral Nutr* 1997;21:353–6.
119. Auvin S, Michaud L, Guimber D, et al. Percutaneous endoscopic jejunostomy for decompression in an infant with short-bowel syndrome. *Endoscopy* 2002;34:240.
120. Silk DB. The evolving role of post-ligament of Trietz nasojejunal feeding in enteral nutrition and the need for improved feeding tube design and placement methods. *J Parenter Enteral Nutr* 2011;35:303–7.
121. Lopez-Herce J. Gastrointestinal complications in critically ill patients: what differs between adults and children? *Curr Opin Clin Nutr Metab Care* 2009;12:180–5.
122. Dellagrammaticas HD, Duerden BI, Milner RD. Upper intestinal bacterial flora during transpyloric feeding. *Arch Dis Child* 1983;58:115–9.
123. Bechtold ML, Nguyen DL, Palmer LB, et al. Nasal bridles for securing nasogastric tubes: a meta-analysis. *Nutr Clin Pract* 2014;29:667–71.
124. Ltd. GE. AMT Bridle™ Pocket Guide. <http://www.gbukenteral.com/pdf/Bridge-insertion-pocket-guide.pdf>. Accessed January 2, 2018.

125. NICE. Infection: Prevention and Control of Healthcare-Associated Infections in Primary and Community Care. <https://www.nice.org.uk/guidance/cg139/evidence/control-full-guideline-185186701>. Accessed January 2, 2018.
126. NPSA. Patient Safety Alert: Reducing the Harm Caused by Misplaced Nasogastric Feeding Tubes in Adults, Children and Infants. <http://www.nrls.npsa.nhs.uk/resources/type/alerts/?entryid45=129640>. Accessed January 2, 2018.
127. Macqueen S, Bruce E, Gibson F. The Great Ormond Street Hospital Manual of Children's Nursing Practices. London: Wiley-Blackwell; 2005.
128. Dandeles LM, Lodolce AE. Efficacy of agents to prevent and treat enteral feeding tube clogs. *Ann Pharmacother* 2011;45:676–80.
129. Forsythe SJ. Enterobacter sakazakii and other bacteria in powdered infant milk formula. *Matern Child Nutr* 2005;1:44–50.
130. WHO. Safe Preparation, Storage and Handling of Powdered Infant Formula Guidelines. http://www.who.int/foodsafety/publications/micro/pif_guidelines.pdf. Accessed January 2, 2018.
131. De Leeuw I, Van Alsenoy L. Bacterial contamination of the feeding bag during catheter jejunostomy: exogenous or endogenous Origin? *JPEN J Parenter Enteral Nutr* 1984;8:591–2.
132. Taylor LJ, Faria SH. Caring for the patient with a gastrostomy/jejunostomy tube. *Home Care Provid* 1997;2:221–4.
133. Young CK, White S. Preparing patients for tube feeding at home. *Am J Nurs* 1992;92:46–53.
134. MDA. Medical Devices Agency. Safety Notice SN2000(27). Enteral Feeding Systems. 2000.
135. Khair J. Managing home enteral tube feeding for children. *Br J Community Nurs* 2003;8:116–26.
136. Lowry S, Johnston SD. Who follows up patients after PEG tube insertion? *Ulster Med J* 2007;76:88–90.
137. Madigan SM. Home enteral-tube feeding: the changing role of the dietitian. *Proc Nutr Soc* 2003;62:761–3.
138. Agostoni C, Axelson I, Colomb V, et al. The need for nutrition support teams in pediatric units: a commentary by the ESPGHAN committee on nutrition. *J Pediatr Gastroenterol Nutr* 2005;41:8–11.
139. Krom H, De Winter JP, Kindermann A. Development, prevention, and treatment of feeding tube dependency. *Eur J Pediatr* 2017;176:683–8.
140. Wilken M, Cremer V, Berry J, et al. Rapid home-based weaning of small children with feeding tube dependency: positive effects on feeding behaviour without deceleration of growth. *Arch Dis Child* 2013;98:856–61.
141. Gottrand F, Sullivan PB. Gastrostomy tube feeding: when to start, what to feed and how to stop. *Eur J Clin Nutr* 2010;64(suppl 1):S17–21.
142. Kindermann A, Kneepkens CM, Stok A, et al. Discontinuation of tube feeding in young children by hunger provocation. *J Pediatr Gastroenterol Nutr* 2008;47:87–91.
143. Mahant S, Jovcevska V, Cohen E. Decision-making around gastrostomy-feeding in children with neurologic disabilities. *Pediatrics* 2011;127:e1471–8.
144. Craig GM, Carr LJ, Cass H, et al. Medical, surgical, and health outcomes of gastrostomy feeding. *Dev Med Child Neurol* 2006;48:353–60.
145. Korner U, Bondolfi A, Buhler E, et al. Ethical and legal aspects of enteral nutrition. *Clin Nutr* 2006;25:196–202.
146. Majka AJ, Wang Z, Schmitz KR, et al. Care coordination to enhance management of long-term enteral tube feeding: a systematic review and meta-analysis. *JPEN J Parenter Enteral Nutr* 2014;38:40–52.
147. Klek S, Szybinski P, Sierzega M, et al. Commercial enteral formulas and nutrition support teams improve the outcome of home enteral tube feeding. *JPEN J Parenter Enteral Nutr* 2011;35:380–5.
148. Best C, Hitchings H. Enteral tube feeding—from hospital to home. *Br J Nurs* 2010;19:76–9.
149. De Luis DA, Izaola O, Cuellar LA, et al. Experience over 12 years with home enteral nutrition in a healthcare area of Spain. *J Hum Nutr Diet* 2013;26(suppl 1):39–44.
150. Silver HJ, Wellman NS, Arnold DJ, et al. Older adults receiving home enteral nutrition: enteral regimen, provider involvement, and health care outcomes. *JPEN J Parenter Enteral Nutr* 2004;28:92–8.